

学位論文の要旨

氏名 小池 抄代

学位論文名 Advanced Glycation End-Products Induce Apoptosis of Vascular Smooth Muscle Cells: A Mechanism for Vascular Calcification
発表雑誌名 International Journal of Molecular Sciences
(巻, 初頁～終頁, 年) (17, 1567, 2016)
著者名 Sayo Koike, Shozo Yano, Sayuri Tanaka, Abdullah M. Sheikh, Atsushi Nagai and Toshitsugu Sugimoto

論文内容の要旨

INTRODUCTION

Vascular complication is an important aspect of the pathological course of diabetes mellitus, and affects the disease-related morbidity and mortality. For the development of such complications, hyperglycemia is suggested to play a central role. Hence, in diabetic patients, a long term intensive control of glycemic status is associated with a significantly reduced risk of micro- and macro-vascular complications, which persists even in good glycemic control afterwards. Advanced glycation end-products (AGEs), which are generated by reducing sugars in non-enzymatic reaction with proteins (Maillard reaction), well explain such a “legacy” effect owing to its difficulty in degradation and clearance from the diabetic tissue.

Mönckeberg-type arterial calcification in the media, which is a characteristic feature of patients with diabetes as well as chronic kidney disease (CKD), develops and progresses time dependently. As a mechanism of vascular calcification, vascular smooth muscle cells (VSMCs) apoptosis is reported to be important. Previous studies have reported that AGEs suppressed cellular differentiation and maturation, and induced apoptosis in many types of cells. However, the effects of AGEs on VSMCs apoptosis and its relationship with vascular calcification are largely unknown. Previously, we have demonstrated that glycolaldehyde-derived AGEs induce calcium deposition in rat VSMCs through excessive generation of reactive oxygen species (ROS) and phenotypic transition into osteoblast like cells. In this study, we aimed to investigate the role of glycolaldehyde-derived AGEs on VSMCs apoptosis, with emphasis on the underlying mechanism.

MATERIALS AND METHODS

A7r5 cells (rat aortic VSMC line) were cultured in Dulbecco’s modified Eagle’s medium (DMEM) containing 10 % FBS at 37 °C in a fully humidified atmosphere of 5 % CO₂ in air. AGE3-bovine serum albumin (BSA) was prepared by incubating BSA with glycolaldehyde and 5 mM diethylenetriamine pentaacetic acid (DTPA) in 0.2 M phosphate buffer (pH 7.4) at 37 °C for 7 days.

For induction of calcification, growth medium was switched to calcification medium (DMEM containing 10% FBS, 10 mM sodium pyruvate, 10^{-7} M insulin, 100 U/mL penicillin, 100 mg/mL streptomycin, and β -glycerophosphate) after reaching the confluency. The cells were treated with 100 μ g/mL of AGE3-BSA or the control BSA (cBSA). Calcium deposition was determined colorimetrically by *O*-cresolphthalein complexone method (calcium C-test Wako) on day3. Apoptotic cells were identified by Terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) staining. Apoptosis was semi-quantitatively evaluated by the ratio of TUNEL positive cell number divided by Hoechst positive cell number. To quantify apoptosis, an enzyme-linked immunosorbent assay (ELISA) for histone-complexed DNA fragments was also employed. Real-time PCR was performed to determine the mRNA levels for Nox1, Nox4 and p22^{phox} using the QuantiTect SYBR PCR kit. RNA interference technique was used to down-regulate the expression of Nox4 and p22^{phox} in A7r5 cells. SMARTpool small interfering RNA (siRNA) and the reagents for these genes were designed and synthesized by Dharmacon. The expression at the protein level was determined by immunofluorescence and quantified using ImageJ software (<http://imagej.net/ImageJ>). Dihydroethidium (DHE) assay was performed to determine cellular ROS levels.

Statistical evaluation of the differences between the groups was carried out with unpaired *t*-test and/or one-way analysis of variance (ANOVA) followed by Fisher's protected least significant difference.

RESULTS AND DISCUSSION

To examine effects of apoptosis on calcium deposition, A7r5 cells were treated with general caspase inhibitor Z-VAD-FMK (10 μ M) or the control Z-FA-FMK (10 μ M) for three days. AGE3-BSA-induced calcium deposition was significantly inhibited by the treatment with caspase inhibitor. This suggests that AGE-induced calcium deposition is mediated by apoptotic cell death in VSMCs. Thus, we investigated AGE-induced apoptosis and the mechanism in A7r5 cells.

The cells were treated with cBSA or increasing concentration of AGE3-BSA (25, 50, 100, 200, and 300 μ g/mL) with calcification medium. On day 5, apoptotic cell death was measured using an ELISA-based method. Up to 50 μ g/mL concentration, AGE3-BSA did not affect A7r5 apoptosis. In contrast, AGE3-BSA significantly increased apoptosis from 100 μ g/mL concentration. However, we did not find any dose-dependent effect of AGE3-BSA beyond 100 μ g/mL concentration. As AGE3-BSA showed maximum apoptotic effect at 100 μ g/mL concentration, we used this dose in all subsequent experiments.

Analysis of apoptosis by TUNEL assay showed that AGE3-BSA markedly increased the number of TUNEL positive cells. Interestingly, pretreatment of cells with NAD(P)H oxidase inhibitor including GKT137831 (20 μ M) or VAS2870 (10 μ M), markedly decreased the number of TUNEL positive cells. Quantification analysis also showed that the percentage of TUNEL positive cells in a total cell culture population was significantly increased by AGE3-BSA treatment, and such effect of

AGE3-BSA was greatly inhibited by NAD(P)H oxidase inhibitors. These findings suggest that AGE3-BSA-induced apoptosis of VSMC was mediated by the activation of NAD(P)H oxidase.

To evaluate further the roles of NAD(P)H oxidase in AGE3-induced apoptosis of VSMCs, we checked the effects of AGE3-BSA on the mRNA and protein expression of the components of NAD(P)H oxidase. Three days after incubation with AGE3-BSA or cBSA, total RNA was isolated from A7r5 cells, and the mRNA expression was assessed by real-time PCR. The results showed that AGE3-BSA treatment significantly increased the expression of Nox1, Nox4 and p22^{phox} mRNA. These findings were consistent with the protein expression of Nox4 and p22^{phox}. Next, we checked whether such increased production of Nox4 and p22^{phox} proteins by AGE3-BSA has any functional significance. In DHE assay, AGE3-BSA significantly increased cellular ROS level compared to medium treated and cBSA treated conditions.

Next, we examined effects of the silencing of Nox4 and p22^{phox} on AGE-induced apoptosis. The real-time PCR results showed that both Nox4 and p22^{phox} mRNA levels were decreased to 5%–20% after mRNA specific siRNA transfection, indicating their sufficient silencing effect. Importantly, AGE3-BSA-induced A7r5 apoptosis was markedly inhibited (42% or 47%) by transfection of either Nox4 or p22^{phox} siRNA, compared to control (scramble siRNA transfection). Double knockdown of Nox4 and p22^{phox} showed a similar inhibitory effect on apoptosis as single gene silencing (42%).

The present study demonstrated that NAD(P)H oxidase-derived oxidative stress is involved in AGEs-induced apoptosis of VSMCs. This might be important to understand the pathogenesis of vascular calcification in diabetes and CKD. Since excessive ROS is thought to stimulate the production of AGEs, the generation of AGEs and ROS might activate a positive feedback loop through NF- κ B as well as receptor for AGE (RAGE), leading to the development of cardiovascular diseases.


In vasculature, ROS is generated mainly by NAD(P)H oxidase, which is composed of Nox isoforms, p22^{phox}, and associated proteins such as p47^{phox} as a subunit. As previous studies demonstrated that the activity of p22^{phox} as well as Nox4 is associated with their mRNA levels, increased expression of Nox4 or p22^{phox} is most probably responsible for ROS generation. However, we did not find any synergistic or additive effect of double silencing of Nox4 and p22^{phox} on the apoptosis. This result indicates that the presence of both Nox4 and p22^{phox} is essential for functional activity of NAD(P)H oxidase. Moreover, the function of one of them cannot compensate the other components. Since silencing of Nox1 did not affect AGE-induced calcium deposition in our previous study, we speculate that expression level of Nox1 is of little importance for calcium deposition in VSMCs. Future studies need to address the roles of Nox1 and p47^{phox} in VSMCs.

CONCLUSION

AGEs stimulate VSMCs apoptosis through excessive ROS generation. Component of NAD(P)H oxidase such as Nox4 may be a good candidate of new strategy to prevent vascular calcification.

論文審査及び最終試験又は学力の確認の結果の要旨

①・乙	氏名	小池 抄代
学位論文名	Advanced Glycation End-Products Induce Apoptosis of Vascular Smooth Muscle Cells: A Mechanism for Vascular Calcification	
学位論文審査委員	主査	竹谷 健
	副査	木下 芳一
	副査	丸山 理留敬



論文審査の結果の要旨

糖尿病患者の予後の中心的役割を果たしている心血管合併症において、タンパク質の糖化反応によって生成される終末糖化産物 (AGE) が関与している可能性が示唆されている。また、血管の石灰化、特に中膜の石灰化が、糖尿病や慢性腎臓病患者において心血管死と関連することが近年報告された。申請者らは、これまでAGEが酸化ストレスを介して血管平滑筋細胞を骨芽細胞様細胞に形質転換させ、カルシウム (Ca) 沈着を来すことで血管の石灰化に寄与していることを明らかにした。このAGEはアポトーシスも誘導するため、今回、AGEが血管平滑筋細胞のアポトーシスをもたらし、中膜の石灰化に関与するかどうかを検討した。ラットの大動脈平滑筋細胞 (A7r5) とグリコールアルデヒドから作成したAGE3を用いて検討した結果、AGE3はA7r5細胞のCa沈着を有意に促進したが、caspase 阻害薬 (Z-VAD-FMK) の添加により抑制されたことから、AGE3誘導性のCa沈着はアポトーシスを介していることが示唆された。また、DNA断片化を検出するELISA法によりアポトーシスを定量的に調べた結果、AGE3は100 µg/mLからアポトーシスを誘導し、TUNEL法でもAGE3によりアポトーシスの促進を認めた。ここで、AGEが酸化ストレスに関与することから活性酸素を産生するNAD(P)H oxidaseの阻害薬 (GKT 137831、VAS 2870) を用いたところ、AGE3によるアポトーシスが抑制されたことからAGE誘導性のアポトーシスにはNAD(P)H oxidaseの活性化が関与していることが明らかになった。さらに、NAD(P)H oxidase の構成分子であるNox4とp22^{phox}の発現がmRNAとタンパクの両方においてAGE3により上昇しており、Reactive oxygen speciesレベルの増加もみられた。そこでsiRNAを用いてNox4とp22^{phox}をそれぞれ単独、もしくは同時にノックダウンしたところ、AGE3誘導性のアポトーシスは有意に、且ついずれも同程度に抑制された。以上の結果より、AGE3は酸化ストレスを介して血管平滑筋細胞のアポトーシスを誘導し、Ca沈着を促進することが示されたため、AGEシグナルやNAD(P)H oxidaseは、血管石灰化・動脈硬化の治療標的となる可能性がある。

最終試験又は学力の確認の結果の要旨

申請者は糖尿病患者の予後に関して重要な役割を果たす心血管合併症の原因の1つである中膜の石灰化が、糖尿病で増加する終末糖化産物 (AGE) が活性酸素を介して血管平滑筋をアポトーシスさせることで引き起こされることを明らかにした。これは血管の石灰化だけでなく動脈硬化の病態解明や治療法の開発にも貢献しうる研究で、関連領域の知識も豊富であることより、学位授与に値すると判断した。
(主査 竹谷健)

申請者は大血管の培養平滑筋細胞を用いて、糖尿病で増加する終末糖化産物が平滑筋細胞のアポトーシスを介してカルシウム沈着を促進することを明らかとした。周辺領域の知識も豊富であり博士の学位に値すると判定した。
(副査 木下芳一)

申請者はAGE3が酸化ストレスを介して血管中膜平滑筋のアポトーシスを引き起こし、さらに石灰化を促進することを証明した。本研究はこの経路が治療標的となる可能性を示唆したもので、臨床的に意義あるものと考えられる。本審査における質疑応答は的確であり周辺知識も豊富であったため、学位授与に値すると判断した。
(副査 丸山理留敬)

(備考) 要旨は、それぞれ400字程度とする。