



Bosonic seesaw mechanism in a classically conformal extension of the Standard Model



Naoyuki Haba^a, Hiroyuki Ishida^a, Nobuchika Okada^b, Yuya Yamaguchi^{a,c,*}

^a Graduate School of Science and Engineering, Shimane University, Matsue 690-8504, Japan

^b Department of Physics and Astronomy, University of Alabama, Tuscaloosa, AL 35487, USA

^c Department of Physics, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan

ARTICLE INFO

Article history:

Received 1 December 2015

Received in revised form 20 January 2016

Accepted 25 January 2016

Available online 29 January 2016

Editor: J. Hisano

ABSTRACT

We suggest the so-called bosonic seesaw mechanism in the context of a classically conformal $U(1)_{B-L}$ extension of the Standard Model with two Higgs doublet fields. The $U(1)_{B-L}$ symmetry is radiatively broken via the Coleman–Weinberg mechanism, which also generates the mass terms for the two Higgs doublets through quartic Higgs couplings. Their masses are all positive but, nevertheless, the electroweak symmetry breaking is realized by the bosonic seesaw mechanism. Analyzing the renormalization group evolutions for all model couplings, we find that a large hierarchy among the quartic Higgs couplings, which is crucial for the bosonic seesaw mechanism to work, is dramatically reduced toward high energies. Therefore, the bosonic seesaw is naturally realized with only a mild hierarchy, if some fundamental theory, which provides the origin of the classically conformal invariance, completes our model at some high energy, for example, the Planck scale. We identify the regions of model parameters which satisfy the perturbativity of the running couplings and the electroweak vacuum stability as well as the naturalness of the electroweak scale.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

In the Standard Model (SM), the electroweak symmetry breaking is realized by the negative mass term in the Higgs potential, which seems to be artificial because there is nothing to stabilize the electroweak scale. If new physics takes place at a very high energy, e.g. the Planck scale, the mass term receives large corrections which are quadratically sensitive to the new physics scale, so that the electroweak scale is not stable against the corrections. This is the so-called gauge hierarchy problem. It is well known that supersymmetry (SUSY) can solve this problem. Since the mass corrections are completely canceled by the SUSY partners, no fine-tuning is necessary to reproduce the electroweak scale correctly, unless the SUSY breaking scale is much higher than the electroweak scale. On the other hand, since no indication of SUSY particles has been obtained in the large hadron collider (LHC) experiments, one may consider other solutions to the gauge hierarchy problem without SUSY.

In this direction, recently a lot of works have been done in models based on a classically conformal symmetry. There are $U(1)$ gauge extension [1–23], and non-Abelian gauge extension,

in which conformal symmetry is broken by radiative corrections [15,24–28] and strong dynamics [29–37]. In addition, there are also non-gauge extended models [see Ref. [38] and therein].¹ This direction is based on the argument by Bardeen [39] that the quadratic divergence in the Higgs mass corrections can be subtracted by a boundary condition of some ultraviolet complete theory, which is classically conformal, and only logarithmic divergences should be considered (see Ref. [6] for more detailed discussions). If this is the case, imposing the classically conformal symmetry to the theory is another way to solve the gauge hierarchy problem. Since there is no dimensionful parameter in this class of models, the classically conformal symmetry must be broken by quantum corrections. This structure fits the model first proposed by Coleman and Weinberg [40], where a model is defined as a massless theory and the classically conformal symmetry is radiatively broken by the Coleman–Weinberg (CW) mechanism, generating a mass scale through the dimensional transmutation.

* Corresponding author at: Department of Physics, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan.

E-mail address: yy@particle.sci.hokudai.ac.jp (Y. Yamaguchi).

¹ In Ref. [38], the upper bound on the mass of the lightest additional scalar boson is obtained as $\simeq 543$ GeV, which is independent of its isospin and hypercharge. Thus, the classically conformal model is strongly constrained without gauge extension.

In this paper we propose a classically conformal $U(1)_{B-L}$ extended SM with two Higgs doublets. An SM singlet, $B-L$ Higgs field develops its vacuum expectation value (VEV) by the CW mechanism, and the $U(1)_{B-L}$ symmetry is radiatively broken. This gauge symmetry breaking also generates the mass terms for the two Higgs doublets through quartic couplings between the two Higgs doublets and the $B-L$ Higgs field. We assume the quartic couplings to be all positive at the $U(1)_{B-L}$ breaking scale but, nevertheless, the electroweak symmetry breaking is triggered through the so-called bosonic seesaw mechanism [41–43], which is analogous to the seesaw mechanism for the neutrino mass generation and leads to a negative mass squared for the SM-like Higgs doublet. Because a negative quartic coupling may cause vacuum instability, it is important to take all quartic couplings to be positive, while in the conventional models, e.g., Refs. [3] and [29], the mixing coupling between the $SU(2)_L$ doublet and singlet fields is necessarily negative to realize the negative mass term of the SM-like Higgs doublet. Our model guarantees that the mixing couplings are positive at the breaking scale with a hierarchy among the quartic couplings, which successfully derives the bosonic seesaw mechanism. The hierarchy seems to be unnatural, but we find that the renormalization group evolutions of the quartic couplings dramatically reduce the large hierarchy toward high energies. On the other hand, a large hierarchy exists even in the conventional model, that is, the mixing coupling should be much small as $(\text{EW scale})^2/v^2$ with a conformal symmetry breaking scale v , except for $v \sim \mathcal{O}(1)$ TeV. Note that the degree of the hierarchy in our model does not increase as the symmetry breaking scale becomes larger.

In the following, let us explain our model in detail. We consider an extension of the SM with an additional $U(1)_{B-L}$ gauge symmetry. Our model has three scalar fields, that is, two Higgs doublets (H_1 and H_2) and one SM singlet, $B-L$ Higgs field (Φ) are introduced. The $U(1)_{B-L}$ charges of H_1 , H_2 , and Φ are 0, 4, and 2, respectively. As is well known, the introduction of the three right-handed neutrinos (N^i , $i = 1, 2, 3$) with a $U(1)_{B-L}$ charge is crucial to make the model free from all the gauge and gravitational anomalies. In addition, we impose a classically conformal symmetry to the model, under which the scalar potential is given by

$$V = \lambda_1 |H_1|^4 + \lambda_2 |H_2|^4 + \lambda_3 |H_1|^2 |H_2|^2 + \lambda_4 (H_2^\dagger H_1)(H_1^\dagger H_2) + \lambda_\Phi |\Phi|^4 + \lambda_{H_1\Phi} |H_1|^2 |\Phi|^2 + \lambda_{H_2\Phi} |H_2|^2 |\Phi|^2 + (\lambda_{\text{mix}} H_2^\dagger H_1) \Phi^2 + h.c.). \quad (1)$$

Here, all of the dimensionful parameters are prohibited by the classically conformal symmetry. In this system, the $U(1)_{B-L}$ symmetry must be radiatively broken by quantum effects, i.e., the CW mechanism. The CW potential for Φ is described as

$$V_\Phi(\phi) = \frac{1}{4} \lambda_\Phi (v_\Phi) \phi^4 + \frac{1}{8} \beta_{\lambda_\Phi} (v_\Phi) \phi^4 \left(\ln \frac{\phi^2}{v_\Phi^2} - \frac{25}{6} \right), \quad (2)$$

where $\Re[\Phi] = \phi/\sqrt{2}$, and $v_\Phi = \langle \phi \rangle$ is the VEV of Φ . When the beta function β_{λ_Φ} is dominated by the $U(1)_{B-L}$ gauge coupling (g_{B-L}) and the Majorana Yukawa couplings of right-handed neutrinos (Y_M), the minimization condition of V_Φ approximately leads to

$$\lambda_\Phi \simeq \frac{11}{6\pi^2} (6g_{B-L}^4 - \text{tr} Y_M^4), \quad (3)$$

where all parameters are evaluated at v_Φ . Through the $U(1)_{B-L}$ symmetry breaking, the mass terms of the two Higgs doublets arise from the mixing terms between $H_{1,2}$ and Φ , and the scalar mass squared matrix is read as

$$-\mathcal{L} = \frac{1}{2} (H_1, H_2) \begin{pmatrix} \lambda_{H_1\Phi} v_\Phi^2 & \lambda_{\text{mix}} v_\Phi^2 \\ \lambda_{\text{mix}} v_\Phi^2 & \lambda_{H_2\Phi} v_\Phi^2 \end{pmatrix} \begin{pmatrix} H_1 \\ H_2 \end{pmatrix} \approx \frac{1}{2} (H'_1, H'_2) \begin{pmatrix} \lambda_{H_1\Phi} v_\Phi^2 - \frac{\lambda_{\text{mix}}^2 v_\Phi^2}{\lambda_{H_2\Phi}} & 0 \\ 0 & \lambda_{H_2\Phi} v_\Phi^2 \end{pmatrix} \begin{pmatrix} H'_1 \\ H'_2 \end{pmatrix}, \quad (4)$$

where H'_1 and H'_2 are mass eigenstates, and we have assumed a hierarchy among the quartic couplings as $0 \leq \lambda_{H_1\Phi} \ll \lambda_{\text{mix}} \ll \lambda_{H_2\Phi}$ at the scale $\mu = v_\Phi$. In the next section, we will show that this hierarchy is dramatically reduced toward high energies in their renormalization group evolutions. Because of this hierarchy, mass eigenstates H'_1 and H'_2 are almost composed of H_1 and H_2 , respectively. Hence, we approximately identify H'_1 with the SM-like Higgs doublet. Note that even though all quartic couplings are positive, the SM-like Higgs doublet obtains a negative mass squared for $\lambda_{H_1\Phi} \ll \lambda_{\text{mix}}^2/\lambda_{H_2\Phi}$, and hence the electroweak symmetry is broken. This is the so-called bosonic seesaw mechanism [41–43].

In more precise analysis for the electroweak symmetry breaking, we take into account a scalar one-loop diagram through the quartic couplings, λ_3 and λ_4 , and the SM-like Higgs doublet mass is given by

$$m_h^2 \simeq \lambda_{H_2\Phi} v_\Phi^2 \left[\frac{1}{2} \left(\frac{\lambda_{\text{mix}}}{\lambda_{H_2\Phi}} \right)^2 + \frac{2\lambda_3 + \lambda_4}{16\pi^2} \right], \quad (5)$$

where we have omitted the $\lambda_{H_1\Phi}$ term in the second line, and the observed Higgs boson mass $M_h = 125$ GeV is given by $M_h = m_h/\sqrt{2}$.

In addition to the scalar one-loop diagram, one may consider other Higgs mass corrections coming from a neutrino one-loop diagram and two-loop diagrams involving the $U(1)_{B-L}$ gauge boson (Z') and the top Yukawa coupling, which are, respectively, found to be [3]

$$\delta m_h^2 \sim \frac{Y_\nu^2 Y_M^2 v_\Phi^2}{16\pi^2}, \quad \delta m_h^2 \sim \frac{Y_t^2 g_{B-L}^4 v_\Phi^2}{(16\pi^2)^2}, \quad (6)$$

where Y_ν and y_t are Dirac Yukawa couplings of neutrino and top quark, respectively. It turns out that these contributions are negligibly small compared to the scalar one-loop correction in Eq. (5). As we will discuss in the next section, the quartic couplings λ_3 and λ_4 should be sizable $\lambda_{3,4} \gtrsim 0.15$ in order to stabilize the electroweak vacuum. The neutrino one-loop correction is roughly proportional to the active neutrino mass by using the seesaw relation, and it is highly suppressed by the lightness of the neutrino mass. The two-loop corrections with the Z' boson are suppressed by a two-loop factor $1/(16\pi^2)^2$. Unless g_{B-L} is large, the two-loop corrections are smaller than the scalar one-loop correction.

The other scalar masses are approximately given by

$$M_\phi^2 = \frac{6}{11} \lambda_\Phi v_\Phi^2, \quad (7)$$

$$M_H^2 = M_A^2 = \lambda_{H_2\Phi} v_\Phi^2 + (\lambda_3 + \lambda_4) v_H^2, \quad (8)$$

$$M_{H^\pm}^2 = \lambda_{H_2\Phi} v_\Phi^2 + \lambda_3 v_H^2, \quad (9)$$

where M_ϕ is the mass of the SM singlet scalar, M_H (M_A) is the mass of CP-even (CP-odd) neutral Higgs boson, and M_{H^\pm} is the mass of charged Higgs boson. The extra heavy Higgs bosons are almost degenerate in mass. The masses of the Z' boson and the right-handed neutrinos are given by

$$M_{Z'} = 2g_{B-L} v_\Phi, \quad (10)$$

$$M_N = \sqrt{2} y_M v_\Phi \simeq \left[\frac{3}{2N_\nu} \left(1 - \frac{\pi^2 \lambda_\Phi}{11g_{B-L}^4} \right) \right]^{1/4} M_{Z'}, \quad (11)$$




Fig. 1. Renormalization group evolutions of the quartic couplings for $v_\phi = 10$ TeV (solid) and 100 TeV (dashed). The red, green, and blue lines correspond to $\lambda_{H1\phi}$, $\lambda_{H2\phi}$ and λ_{mix} , respectively. The rightmost vertical line shows the reduced Planck scale. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

where we have used $\text{tr}Y_M = N_\nu y_M$, for simplicity, and N_ν stands for the number of relevant Majorana couplings. In the following analysis, we will take $N_\nu = 1$ for simplicity, because our final results are almost insensitive to N_ν . In the last equality in Eq. (11), we have used Eq. (3).

Before presenting our numerical results, we first discuss constraints on the model parameters from the perturbativity and the stability of the electroweak vacuum in the renormalization group evolutions. In our analysis, all values of couplings are given at $\mu = v_\phi$. For v_ϕ at the TeV scale, we find the constraint $g_{B-L} \lesssim 0.3$ to avoid the Landau pole of the gauge coupling below the Planck scale, while a more severe constraint $g_{B-L} \lesssim 0.2$ is obtained to avoid a blowup of the quartic coupling λ_2 below the Planck scale. From $g_{B-L} \lesssim 0.2$ and the experimental bound $M_{Z'} > 2.9$ TeV on the Z' boson mass [44,45], we find $v_\phi > 7.25$ TeV. The electroweak vacuum stability, in other words, $\lambda_H(\mu) > 0$ for any scales between the electroweak scale and the Planck scale, can be realized by sufficiently large λ_3 and/or λ_4 as $\lambda_3 = \lambda_4 \gtrsim 0.15$. To keep their perturbativity below the Planck scale, $\lambda_3 = \lambda_4 \lesssim 0.48$ must be satisfied, while we will find that the naturalness of the electroweak scale leads to a more severe upper bound.

To realize the hierarchy $\lambda_{H1\phi} \ll \lambda_{\text{mix}} \ll \lambda_{H2\phi}$, we take $\lambda_{H1\phi} = 0$, for simplicity. When we consider λ_{mix} in the range of $0 < \lambda_{\text{mix}} < 0.1 \times \lambda_{H2\phi}$, the relation between v_ϕ and $\lambda_{H2\phi}$ obtained by Eq. (5) is almost uniquely determined. When we fix $\lambda_3 = \lambda_4 = 0.15$ as an example, we find $1 \text{ TeV} \lesssim \lambda_{H2\phi} v_\phi^2 \lesssim 1.7 \text{ TeV}$ for $v_\phi \gtrsim 10 \text{ TeV}$, which is almost independent of v_ϕ . Since all heavy Higgs boson masses are approximately determined by $\lambda_{H2\phi} v_\phi^2$, they lie in the range between 1 TeV and 1.7 TeV. Such heavy Higgs bosons can be tested at the LHC in the near future.

In Eq. (5), it may be natural for the first term from the tree-level couplings dominates over the second term from the 1-loop correction. This naturalness leads to the constraint of $\lambda_3 = \lambda_4 < 0.26$, which is more severe than the perturbativity bound $\lambda_3 = \lambda_4 \lesssim 0.48$ discussed above. This condition is equivalent to the fact that the origin of the negative mass term mainly comes from the diagonalization of the scalar mass squared matrix in Eq. (4), namely, the bosonic seesaw mechanism.

Now we present the results of our numerical analysis. In Fig. 1, we show the renormalization group evolutions of the quartic couplings. Here, we have taken $\lambda_{H1\phi} = 0$, and $\lambda_{H2\phi} = 10^{-2}$ and 10^{-4} for $v_\phi = 10$ TeV (solid lines) and 100 TeV (dashed lines), respectively. The red, green, and blue lines correspond to the running of $\lambda_{H1\phi}$, $\lambda_{H2\phi}$ and λ_{mix} , respectively. The rightmost vertical line denotes the reduced Planck scale $M_{Pl} = 2.4 \times 10^{18}$ GeV. In this plot, the other input parameters have been set as $g_{B-L} = 0.17$ and $\lambda_3 = \lambda_4 = 0.17$ to realize the electroweak vacuum stability without the

Table 1

Additional vector-like fermions. x is a real number.

| | $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ | $U(1)_{B-L}$ |
|------------|--|--------------|
| $S_{L,R}$ | (1, 1, 0) | x |
| $S'_{L,R}$ | (1, 1, 0) | $x - 2$ |
| $D_{L,R}$ | (1, 2, 1/2) | x |
| $D'_{L,R}$ | (1, 2, 1/2) | $x + 2$ |

Landau pole, and $\lambda_\phi = 10^{-3}$. The value of $\lambda_1 = \lambda_2 = \lambda_H$ at $\mu = v_\phi$ has been evaluated by extrapolating the SM Higgs quartic coupling with $M_h = 125$ GeV from the electroweak scale to v_ϕ . For this parameter choice, the Z' boson and the right-handed neutrinos have the masses of the same order of magnitude as $M_{Z'} = 3.4$ (34) TeV and $M_N = 2.0$ (20) TeV for $v_\phi = 10$ (100) TeV, while the $B - L$ Higgs boson mass is calculated as $M_\phi = 0.23$ (2.3) TeV. As is well-known, $M_\phi \ll M_{Z'}$ is a typical prediction of the CW mechanism. The masses of the heavy Higgs bosons are roughly 1 TeV for both $v_\phi = 10$ TeV and 100 TeV.

In order for the bosonic seesaw mechanism to work, we have assumed the hierarchy among the quartic couplings as $\lambda_{H1\phi} \ll \lambda_{\text{mix}} \ll \lambda_{H2\phi}$ at the scale $\mu = v_\phi$. One may think it unnatural to introduce this large hierarchy by hand. However, we find from Fig. 1 that the large hierarchy between $\lambda_{H1\phi}$ and $\lambda_{H2\phi}$ tends to disappear toward high energies. This is because the beta functions of the small couplings $\beta_{\lambda_{H1\phi}}$ and $\beta_{\lambda_{H2\phi}}$ are not simply proportional to themselves, but include terms given by other sizable couplings, such as $\lambda_3 \lambda_{H2\phi}$ and g_{B-L}^4 . This behavior of reducing the large hierarchy in the renormalization group evolutions is independent of the choice of the boundary conditions for g_{B-L} , λ_3 , λ_4 and λ_ϕ . Therefore, Fig. 1 indicates that once our model is defined at some high energy, say, the Planck scale, the large hierarchy among the quartic couplings, which is crucial for the bosonic seesaw mechanism to work, is naturally achieved from a mild hierarchy at the high energy.

We see in Fig. 1 that λ_{mix} is almost unchanged. This is because $\beta_{\lambda_{\text{mix}}}$ is proportional to λ_{mix} , which is very small. Hence, the hierarchy between λ_{mix} and the other couplings gets enlarged at high energies. To avoid this situation and make our model more natural, one may introduce additional vector-like fermions listed in Table 1, for example. (As another possibility, one may think that some symmetry forbids the λ_{mix} term and it is generated via a small breaking.) Although x is an arbitrary real number, we assume $x \neq 1$ to distinguish the new fermions from the SM leptons. These fermions have Yukawa couplings as

$$\begin{aligned}
 -\mathcal{L}_V = & Y_{SS} \overline{S}_L \Phi S'_R + Y_{SD} \overline{S}_R H_2^\dagger D'_L + Y_{DD} \overline{D}'_L \Phi D_R \\
 & + Y_{DS} \overline{D}_R H_1 S_L + Y'_{SS} \overline{S}_R \Phi S'_L + Y'_{SD} \overline{S}'_L H_2^\dagger D'_R \\
 & + Y'_{DD} \overline{D}'_R \Phi D_L + Y'_{DS} \overline{D}_L H_1 S_R + h.c., \quad (12)
 \end{aligned}$$

so that $\beta_{\lambda_{\text{mix}}}$ includes terms of $Y_{SS} Y_{SD} Y_{DD} Y_{DS}$ and $Y'_{SS} Y'_{SD} \times Y'_{DD} Y'_{DS}$, which are not proportional to λ_{mix} . Accordingly, the minimization condition of V_ϕ is modified to

$$\begin{aligned}
 \lambda_\phi \simeq & \frac{11}{6\pi^2} \left[-\frac{1}{8} \left(Y_{SS}^4 + Y_{SS'}^4 + 2Y_{DD}^4 + 2Y_{DD'}^4 \right) \right. \\
 & \left. + 6g_{B-L}^4 - \text{tr}Y_M^4 \right]. \quad (13)
 \end{aligned}$$

From the conditions $\lambda_\phi > 0$ and $g_{B-L} < 0.2$, the additional Yukawa contribution should satisfy $Y_{SS}^4 + Y_{SS'}^4 + 2Y_{DD}^4 + 2Y_{DD'}^4 \lesssim 3 \times (0.4)^4$. Note that the vector-like fermions masses are dominantly generated by v_ϕ , and they are sufficiently heavy to avoid the current experimental bounds.




Fig. 2. Runnings of quartic couplings for $v_\phi = 100$ TeV with additional vector-like fermions. The vertical axis shows absolute values of quartic couplings. The input parameters are the same as before.

Fig. 2 shows the runnings of the quartic couplings for $v_\phi = 100$ TeV with the additional vector-like fermions. The input parameters are the same as before, while we have taken the Yukawa couplings as $Y_{SS} = Y_{SD} = Y_{DD} = Y_{DS} = 0.2$ and $Y'_{SS} = Y'_{SD} = Y'_{DD} = Y'_{DS} = 0.1$ at $\mu = v_\phi$, for simplicity. Toward high energies, $|\lambda_{\text{mix}}|$ becomes larger, and the hierarchy with the other couplings becomes mild. We can see that $\lambda_{H1\phi}$ is negative below $\mu \simeq 10^8$ GeV,² because the contributions of additional Yukawa couplings to $\beta_{\lambda_{H1\phi}}$ are effective below $\mu \simeq 10^8$ GeV. Above the scale, the contribution of $U(1)_{B-L}$ couplings becomes effective, and then $\lambda_{H1\phi}$ becomes positive. As a result, the large hierarchy at the $U(1)_{B-L}$ symmetry breaking scale can be realized with a mild hierarchy at some high energy. We expect that a ultraviolet complete theory, which provides the origin of the classical conformal invariance, takes place at the high energy.

Acknowledgements

N.O. would like to thank the Particle Physics Theory Group of Shimane University for hospitality during his visit. This work is partially supported by Scientific Grants by the Ministry of Education, Culture, Sports, Science and Technology (Nos. 24540272, 26247038, and 15H01037) and the United States Department of Energy (DE-SC 0013680). The work of Y.Y. is supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists (Grants No. 26-2428).

References

- [1] R. Hempfling, *Phys. Lett. B* 379 (1996) 153, arXiv:hep-ph/9604278.
- [2] W.F. Chang, J.N. Ng, J.M.S. Wu, *Phys. Rev. D* 75 (2007) 115016, arXiv:hep-ph/0701254.
- [3] S. Iso, N. Okada, Y. Orikasa, *Phys. Lett. B* 676 (2009) 81, arXiv:0902.4050 [hep-ph].
- [4] S. Iso, N. Okada, Y. Orikasa, *Phys. Rev. D* 80 (2009) 115007, arXiv:0909.0128 [hep-ph].
- [5] N. Okada, Y. Orikasa, *Phys. Rev. D* 85 (2012) 115006, arXiv:1202.1405 [hep-ph].
- [6] S. Iso, Y. Orikasa, *Prog. Theor. Exp. Phys.* 2013 (2013) 023B08, arXiv:1210.2848 [hep-ph].
- [7] C. Englert, J. Jaeckel, V.V. Khoze, M. Spannowsky, *J. High Energy Phys.* 1304 (2013) 060, arXiv:1301.4224 [hep-ph].
- [8] E.J. Chun, S. Jung, H.M. Lee, *Phys. Lett. B* 725 (2013) 158; E.J. Chun, S. Jung, H.M. Lee, *Phys. Lett. B* 730 (2014) 357, arXiv:1304.5815 [hep-ph].
- [9] I. Oda, *Phys. Lett. B* 724 (2013) 160, arXiv:1305.0884 [hep-ph].
- [10] V.V. Khoze, G. Ro, *J. High Energy Phys.* 1310 (2013) 075, arXiv:1307.3764 [hep-ph].
- [11] M. Hashimoto, S. Iso, Y. Orikasa, *Phys. Rev. D* 89 (1) (2014) 016019, arXiv:1310.4304 [hep-ph].
- [12] M. Lindner, D. Schmidt, A. Watanabe, *Phys. Rev. D* 89 (1) (2014) 013007, arXiv:1310.6582 [hep-ph].
- [13] M. Hashimoto, S. Iso, Y. Orikasa, *Phys. Rev. D* 89 (5) (2014) 056010, arXiv:1401.5944 [hep-ph].
- [14] S. Benic, B. Radovicic, *Phys. Lett. B* 732 (2014) 91, arXiv:1401.8183 [hep-ph].
- [15] V.V. Khoze, C. McCabe, G. Ro, *J. High Energy Phys.* 1408 (2014) 026, arXiv:1403.4953 [hep-ph].
- [16] S. Benic, B. Radovicic, *J. High Energy Phys.* 1501 (2015) 143, arXiv:1409.5776 [hep-ph].
- [17] H. Okada, Y. Orikasa, arXiv:1412.3616 [hep-ph].
- [18] J. Guo, Z. Kang, P. Ko, Y. Orikasa, *Phys. Rev. D* 91 (11) (2015) 115017, arXiv:1502.00508 [hep-ph].
- [19] P. Humbert, M. Lindner, J. Smirnov, *J. High Energy Phys.* 1506 (2015) 035, arXiv:1503.03066 [hep-ph].
- [20] N. Haba, Y. Yamaguchi, arXiv:1504.05669 [hep-ph].
- [21] S. Oda, N. Okada, D.S. Takahashi, arXiv:1504.06291 [hep-ph].
- [22] P. Humbert, M. Lindner, S. Patra, J. Smirnov, arXiv:1505.07453 [hep-ph].
- [23] A.D. Plascencia, arXiv:1507.04996 [hep-ph].
- [24] M. Holthausen, M. Lindner, M.A. Schmidt, *Phys. Rev. D* 82 (2010) 055002, <http://dx.doi.org/10.1103/PhysRevD.82.055002>, arXiv:0911.0710 [hep-ph].
- [25] M. Heikinheimo, A. Racioppi, M. Raidal, C. Spethmann, K. Tuominen, *Mod. Phys. Lett. A* 29 (2014) 1450077, <http://dx.doi.org/10.1142/S0217732314500771>, arXiv:1304.7006 [hep-ph].
- [26] M. Heikinheimo, A. Racioppi, M. Raidal, C. Spethmann, K. Tuominen, *Nucl. Phys. B* 876 (2013) 201, <http://dx.doi.org/10.1016/j.nuclphysb.2013.08.006>, arXiv:1305.4182 [hep-ph].
- [27] T. Hambye, A. Strumia, *Phys. Rev. D* 88 (2013) 055022, <http://dx.doi.org/10.1103/PhysRevD.88.055022>, arXiv:1306.2329 [hep-ph].
- [28] C.D. Carone, R. Ramos, *Phys. Rev. D* 88 (2013) 055020, <http://dx.doi.org/10.1103/PhysRevD.88.055020>, arXiv:1307.8428 [hep-ph].
- [29] T. Hur, P. Ko, *Phys. Rev. Lett.* 106 (2011) 141802, <http://dx.doi.org/10.1103/PhysRevLett.106.141802>, arXiv:1103.2571 [hep-ph].
- [30] M. Holthausen, J. Kubo, K.S. Lim, M. Lindner, *J. High Energy Phys.* 1312 (2013) 076, [http://dx.doi.org/10.1007/JHEP12\(2013\)076](http://dx.doi.org/10.1007/JHEP12(2013)076), arXiv:1310.4423 [hep-ph].
- [31] J. Kubo, K.S. Lim, M. Lindner, *Phys. Rev. Lett.* 113 (2014) 091604, <http://dx.doi.org/10.1103/PhysRevLett.113.091604>, arXiv:1403.4262 [hep-ph].
- [32] J. Kubo, K.S. Lim, M. Lindner, *J. High Energy Phys.* 1409 (2014) 016, [http://dx.doi.org/10.1007/JHEP09\(2014\)016](http://dx.doi.org/10.1007/JHEP09(2014)016), arXiv:1405.1052 [hep-ph].
- [33] M. Heikinheimo, C. Spethmann, *J. High Energy Phys.* 1412 (2014) 084, [http://dx.doi.org/10.1007/JHEP12\(2014\)084](http://dx.doi.org/10.1007/JHEP12(2014)084), arXiv:1410.4842 [hep-ph].
- [34] Y. Ametani, M. Aoki, H. Goto, J. Kubo, *Phys. Rev. D* 91 (11) (2015) 115007, <http://dx.doi.org/10.1103/PhysRevD.91.115007>, arXiv:1505.00128 [hep-ph].
- [35] J. Kubo, M. Yamada, arXiv:1505.05971 [hep-ph].
- [36] C.D. Carone, R. Ramos, *Phys. Lett. B* 746 (2015) 424, <http://dx.doi.org/10.1016/j.physletb.2015.05.044>, arXiv:1505.04448 [hep-ph].
- [37] J. Kubo, M. Yamada, *Prog. Theor. Exp. Phys.* 2015 (9) (2015) 093B01, <http://dx.doi.org/10.1093/ptep/ptv114>, arXiv:1506.06460 [hep-ph].
- [38] K. Hashino, S. Kanemura, Y. Orikasa, *Phys. Lett. B* 752 (2016) 217, <http://dx.doi.org/10.1016/j.physletb.2015.11.044>, arXiv:1508.03245 [hep-ph].
- [39] W.A. Bardeen, FERMILAB-CONF-95-391-T, C95-08-27.3.
- [40] S.R. Coleman, E.J. Weinberg, *Phys. Rev. D* 7 (1973) 1888.
- [41] X. Calmet, *Eur. Phys. J. C* 28 (2003) 451, arXiv:hep-ph/0206091.
- [42] H.D. Kim, *Phys. Rev. D* 72 (2005) 055015, arXiv:hep-ph/0501059.
- [43] N. Haba, N. Kitazawa, N. Okada, *Acta Phys. Pol. B* 40 (2009) 67, arXiv:hep-ph/0504279.
- [44] G. Aad, et al., ATLAS Collaboration, *Phys. Rev. D* 90 (5) (2014) 052005, arXiv:1405.4123 [hep-ex].
- [45] V. Khachatryan, et al., CMS Collaboration, *J. High Energy Phys.* 1504 (2015) 025, arXiv:1412.6302 [hep-ex].

² Although $\lambda_{H1\phi}$ and λ_{mix} become negative, their values can be much smaller than the self-couplings (λ_1 and λ_2). Thus, the vacuum is stable in our model.