



島根大学学術情報リポジトリ
S W A N
Shimane University Web Archives of kNowledge

Title

The protective effect of dietary eicosapentaenoic acid against impairment of spatial cognition learning ability in rats infused with amyloid beta(1-40)

Author(s)

Michio Hashimoto, Shahdat Hossain, Yoko Tanabe, Akiko Kawashima, Tsuyoshi Harada, Takashi Yano, Kiyoshi Mizuguchi, Osamu Shido

Journal

The Journal of nutritional biochemistry, Volume20, Issue12

Published

2009 Dec

URL

<https://doi.org/10.1016/j.jnutbio.2008.08.009>

この論文は出版社版ではありません。
引用の際には出版社版をご確認のうえご利用ください。



The protective effect of dietary eicosapentaenoic acid against impairment of spatial cognition learning ability in rats infused with amyloid $\beta_{(1-40)}$

Michio Hashimoto^{a,*}, Shahdat Hossain^a, Yoko Tanabe^a, Akiko Kawashima^b,
Tsuyoshi Harada^b, Takashi Yano^b, Kiyoshi Mizuguchi^b, Osamu Shido^a

^aDepartment of Environmental Physiology, Shimane University Faculty of Medicine, Izumo, Shimane 693-8501, Japan

^bDevelopmental Research, Pharmaceutical Research Center, Mochida Pharmaceutical Co., Ltd., Gotemba, Shizuoka 412-8524, Japan

Received 22 March 2008; received in revised form 20 August 2008; accepted 25 August 2008

Abstract

Background: Amyloid β ($A\beta$) peptide (1–40) can cause cognitive impairment.

Experimental design: We investigated whether dietary preadministration of eicosapentaenoic acid (EPA) is conducive to cognition learning ability and whether it protects against the impairment of learning ability in rats infused with $A\beta$ peptide (1–40) into the cerebral ventricle.

Results: Dietary EPA administered to rats for 12 weeks before the infusion of $A\beta$ into the rat brain significantly decreased the number of reference memory errors (RMEs) and working memory errors (WMEs), suggesting that chronic administration of EPA improves cognition learning ability in rats. EPA preadministered to the $A\beta$ -infused rats significantly reduced the increase in the number of RMEs and WMEs, with concurrent proportional increases in the levels of corticohippocampal EPA and docosahexaenoic acid (DHA) and in the DHA/arachidonic acid molar ratio. Decrease in oxidative stress in these tissues was evaluated by determining the reactive oxygen species and lipid peroxide levels. cDNA microarray analysis revealed that altered genes included those that control synaptic signal transduction, cell communication, membrane-related vesicular transport functions, and enzymes and several other proteins.

Conclusion: The present study suggests that EPA, by acting as a precursor for DHA, ameliorates learning deficits associated with Alzheimer's disease and that these effects are modulated by the expression of proteins involved in neuronal plasticity.

© 2008 Published by Elsevier Inc.

Keywords: Alzheimer's disease; Hippocampus; Spatial memory; Fatty acid; Rat; Amyloid β

1. Introduction

Fish oil provides a host of health benefits because of its major polyunsaturated fatty acid (PUFA) components: eicosapentaenoic acid [EPA; C20:5(*n*–3)] and docosahexaenoic acid [DHA; C22:6(*n*–3)]. The beneficial effects of fish oil on brain functions, however, have largely focused on and highlighted only DHA, the elongated/desaturated product of EPA. That is probably due to the fact that DHA, but not EPA, constitutes the major PUFA of brain lipids: DHA alone constitutes >17% of the total fatty acids in the rat brain [1], while EPA, as a precursor for DHA, constitutes only a tiny percentage of the total fatty acids in

the brain, thus the extensive studies on the beneficial effects of DHA on cognition learning ability.

Alzheimer's disease (AD) is a primary degenerative disease of the central nervous system, and the histopathological hallmark of AD is the presence of neurofibrillar tangles and amyloid plaques of insoluble amyloid peptide aggregates, which ultimately leads to dementia and behavioral and cognitive impairments [2]. Epidemiological studies show that intake of fish oil is associated with a reduced risk of AD [3,4]. Chronic administration of DHA improves spatial learning ability by increasing the level of DHA in the hippocampus and cerebral cortex of young and aged rats [5,6]. DHA administration also protects against [7] and ameliorates [8] memory deficits in amyloid β ($A\beta$)-peptide-induced AD model rats. DHA protects against behavior deficits and dendritic pathology in the AD mouse model [9].

* Corresponding author. Tel.: +81 853 20 2110; fax: +81 853 20 2110.
E-mail address: michio1@med.shimane-u.ac.jp (M. Hashimoto).




Fig. 1. Schema and schedule of experimental rat groups.

There are few reports on the effects of pure EPA on cognition learning ability. Chronic administration of pure EPA attenuates interleukin 1β , but does not significantly enhance memory in control rats, although whether their normal laboratory chow diet contained any $n-3$ PUFAs is unclear [10]. $n-3$ PUFAs can induce gene expression with concomitant effects on synaptic transmission [11] and related signal transduction. In a parallel set of experiments, we also investigated whether dietary EPA could induce gene expressions related to its beneficial effects on learning-related ability. EPA is probably taken into brain tissue, since the presence of Δ^4 -desaturation enzymes in the brain is still unclear [12]. Nonetheless, ^{14}C -labeled EPA detected in the rat brain 1 h after its oral administration to rats decreases time dependently, while ^{14}C]DHA, a metabolite of EPA, increases time dependently [13]. It is also speculated that in the de novo system, each PUFA that is metabolized after being taken into cerebral endothelial cells and astrocytes (constituent cells of the blood–brain barrier) is released from those cells, and that DHA is taken into neurons as metabolite [14]. Here, we estimated the effects of the chronic administration of pure EPA on spatial learning ability in rats and examined whether EPA can protect against the impairment of learning ability in $\text{A}\beta_{(1-40)}$ -induced AD model rats.

2. Materials and methods

2.1. Animals and diet

The experimental schedule is shown in Fig. 1. Rats were handled and killed in accordance with the procedures outlined in the Guidelines for Animal Experimentation of Shimane Medical University (Shimane, Japan), as compiled from the Guidelines for Animal Experimentation of the Japanese Association for Laboratory of Animal Science. Wistar rats (first generation) (*Jcl*: Wistar; Clea Japan Co., Osaka, Japan) were housed in a room under conditions of

controlled temperature ($23\pm 2^\circ\text{C}$), relative humidity ($50\pm 10\%$) and light–dark cycles (light: 0800–2000 h; dark: 2000–0800 h), and were provided fish-oil-deficient pellet diet (F-1; Funabashi Farm, Funabashi, Japan) and water ad libitum. The inbred third-generation male rats ($n=48$; 5 weeks old) were divided into two groups: an EPA group ($n=24$) administered EPA-95E (300 mg/kg/day; Mochida Pharmaceutical Co., Tokyo, Japan) dissolved in 5% gum arabic solution by gavage for 7 weeks initially, and a control group ($n=21$) given 5% gum arabic solution only.

2.2. Preparation of $\text{A}\beta$ -infused rats

The surgical techniques for preparing $\text{A}\beta$ -infused rats were essentially the same as those described previously [7,8]. Briefly, each rat was anesthetized with sodium pentobarbital (50 mg/kg body weight ip), and its skull was exposed and drilled with two holes (right and left, relative to bregma; 0.8 mm posterior, 1.4 mm lateral) in accordance with the atlas of Paxinos and Watson using a stereotaxic frame (Narishige, Tokyo, Japan). A solvent of 35% (vol/vol) acetonitrile plus 0.1% (vol/vol) trifluoroacetic acid (pH 2.0) was used as vehicle for $\text{A}\beta$ peptide (1–40) (Peptide Inst., Osaka, Japan). A miniosmotic pump (Alzet 2002; Durect Co., Cupertino, CA, USA) containing either $\text{A}\beta$ peptide (1–40) solution ($234\pm 13.9\ \mu\text{l}$) or vehicle alone was quickly implanted into the back of the rats. The outlet of the pump was inserted 3.5 mm into the left ventricle and attached to the skull with screws and dental cement. The infusion rate was $0.56\ \mu\text{l/h}$, and the total amount infused was approximately 4.9–5.5 nmol/L $\text{A}\beta$. Spontaneous infusion for 2 weeks thus brought about completion of the volume used in the miniosmotic pump.

2.3. Radial maze learning ability

Seven weeks after the start of EPA administration, the learning-related behavior of the rats was assessed by their completing a task in an eight-arm radial maze as previously described [5,8]: four reward pellets were placed randomly on four arms of the maze, and the number of total selections in obtaining the four pellets was counted. Two parameters of memory function were examined: reference memory error (RME), which was determined by the number of entries into unbaited arms, and working memory error (WME), which was estimated by the number of repeated entries into arms that had already been visited within a trial. Lower numbers of RMEs and WMEs implied better spatial learning ability in the rats. Performance was calculated from memory-related behavior. Each rat was given two daily trials, 6 days/week for a total of 3 weeks. After completing the behavior test, each of the two groups of rats was subdivided into two groups (allowing for the number of errors made by each rat in the last six trials of the preliminary behavior test) and infused with either $\text{A}\beta$ or the vehicle as follows: a control group was divided into an $\text{A}\beta$ -solvent-infused group [control (vehicle) group; $n=12$] and an $\text{A}\beta$ -infused group ($\text{A}\beta$ group; $n=9$);

143 an EPA group was divided into a vehicle-infused EPA group
144 (EPA+vehicle group; $n=12$) and an A β -infused EPA group
145 (EPA+A β group; $n=12$). The four groups of rats were again
146 behaviorally tested 3 weeks after the implantation of the
147 miniosmotic pump to assess the effect of EPA preadminis-
148 tration on the impairment of learning ability in A β -infused
149 rats. This testing lasted for a total of 3 weeks. The same
150 protocol used for the preliminary behavior test was followed
151 in the final behavior test, except for the adaptation periods.

152 2.4. Preparation of sample

153 After undergoing the behavioral tests for 3 days, the rats
154 were anesthetized with sodium pentobarbital (65 mg/kg
155 body weight, ip), blood was drawn for plasma analysis, and
156 the hippocampus and cerebral cortex were separated as
157 described previously [7]. The tissues were stored at -80°C
158 by flash-freezing in liquid N₂ until use.

159 2.5. Measurement of fatty acid profile and oxidative status

160 The brain samples were immediately homogenized on ice
161 in 1.0 ml of ice-cold 0.32 mol/L sucrose buffer (pH 7.4)
162 containing 2 mmol/L EDTA, 0.5 mg/L leupeptin, 0.5 mg/L
163 pepstatin, 0.5 mg/L aprotinin and 0.2 mmol/L phenylmethyl-
164 sulfonyl fluoride, using a Polytron homogenizer (PCU-2-
165 110; Kinematica GmbH, Steinhofhalde, Switzerland), and
166 the residual tissues were stored at -80°C by flash-freezing in
167 liquid N₂ until use. The homogenates were immediately
168 subjected to the assays described below or stored at -80°C
169 after liquid N₂ flash-freezing and bathing until use.

170 Lipid peroxide (LPO) concentration was assessed by the
171 thiobarbituric-acid-reactive substance assay of Ohkawa et al.
172 [15] as described by Hashimoto et al. [7,8], and its levels
173 were measured in nanomoles of malondialdehyde per
174 milligram of protein. Malondialdehyde levels were calcu-
175 lated relative to a standard preparation of 1,1,3,3-
176 tetraethoxypropane.

177 The levels of reactive oxygen species (ROS) were
178 determined as described previously [7,8]. Briefly, 50 μl of
179 freshly prepared tissue homogenate was mixed with 4.85 ml
180 of 0.1 mol/L potassium phosphate buffer (pH 7.4) and
181 incubated with 2',7'-dichlorofluorescein diacetate (Molecular
182 Probes, Eugene, OR, USA) in methanol at a final
183 concentration of 5 $\mu\text{mol/L}$ for 15 min at 37°C . The dye-
184 loaded samples were centrifuged at $12,500\times g$ for 10 min at
185 4°C . The pellet was mixed on a vortex at 0°C in 5 ml of
186 0.1 mol/L potassium phosphate buffer (pH 7.4) and
187 incubated for 60 min at 37°C . Fluorescence was measured
188 with a Hitachi 850 spectrofluorometer (Hitachi, Tokyo,
189 Japan) at excitation and emission wavelengths of 488 and
190 525 nm, respectively. A cuvette holder was maintained at
191 37°C . ROS was quantified from a dichlorofluorescein
192 standard curve in methanol.

193 The fatty acid compositions of plasma and brain tissues
194 were determined by gas chromatography as described
195 previously [7].

Protein concentration was estimated by the method of
Lowry et al. [16].

2.6. Gene expression analysis

Gene expression analysis was carried out with the
GeneChip system (Affymetrix) in accordance with the
manufacturer's protocol [17]. Briefly, double-stranded DNA
was synthesized from 5 μg of total RNA, and the cDNA
obtained was used as a template for in vitro transcription.
Fragmented in vitro transcripts were hybridized overnight
with Rat Expression Array 230A (Affymetrix), stained,
washed and scanned with an Affymetrix GeneArray
scanner, where the intensity of the fluorescence for each
feature was measured. The expression value (average
difference) for each gene was determined by calculating
the average of differences in intensity (perfect match
intensity minus mismatch intensity) between its probe
pairs. The image files obtained were analyzed with the
Affymetrix data suite system Microarray Suite 5.0 (MAS
5.0). The expression analysis file created from each sample
(chip) was imported into GeneSpring 5.1 (Agilent Tech-
nologies, Inc., Palo Alto, CA) for further data characteriza-
tion. Briefly, a new experiment was generated after
importing data from the same organ in which data were
normalized by array to the 50th percentile of all measure-
ments on that array. Data filtration based on flags present or
marginal in at least one of the samples was first performed,
and a corresponding gene list based on those flags was
generated. Lists of same-phenotype genes that were either
induced or suppressed were created by filtration-on-fold
function. Gene Ontology (GO) category analyses were
performed using the GeneSpring GO browser, which
calculates hypergeometric P -values to measure statistical
significance for a specific GO category.

2.7. Statistical analysis

Results are expressed as mean \pm S.E. Behavioral data were
analyzed by a two-factor (Group and Block) randomized
block factorial analysis of variance (ANOVA), and all other
parameters were analyzed for intergroup differences by one-
way ANOVA. ANOVA was followed by Bonferroni post hoc
comparisons. Correlation was determined by simple regres-
sion analysis. The statistical programs used were GB-STAT
6.5.4 (Dynamic Microsystems, Inc., SilverSpring, MD,
USA) and StatView 4.01 (MindVision Software; Abacus
Concepts, Inc., Berkeley, CA, USA). $P<.05$ was considered
statistically significant.

3. Results

3.1. Body weight

The final body weights did not differ among the groups
(vehicle group: 393 ± 14 ; A β group: 401 ± 12 ; EPA+vehicle
group: 385 ± 36 ; EPA+A β group: 404 ± 35 g). The brain slices




Fig. 2. Effects of the chronic administration of EPA (left) and the infusion of Aβ peptide (1–40) into the rat cerebral ventricle (right) on the number of WMEs evaluated by the radial maze task (see Materials and Methods for details). Left: Control rats (5% gum-arabic-administered rats; $n=21$) and EPA rats ($n=24$). After completing the initial behavior test, each of the two groups (control and EPA groups) was subdivided into two groups and infused with either (right) Aβ (Aβ group; $n=9$) or Aβ solvent (control group; $n=12$): the EPA group was divided into a vehicle-infused EPA group (EPA group; $n=12$) and an Aβ-infused EPA group (EPA+Aβ group; $n=12$). The four groups of rats were again behaviorally tested (with six trials) after the implantation of the miniosmotic pump. Each value represents the number of WMEs, presented as mean±S.E.M. in each block of six trials. The statistical significance of differences between the groups was determined by randomized two-factor (Block and Group) ANOVA followed by Bonferroni post hoc test.

246 prepared 16–17 days after infusion of the Aβ peptides
247 clearly indicated the deposition of the infused Aβ_(1–40) in the
248 corticohippocampal regions (data not shown).




Fig. 3. Effects of the chronic administration of EPA (left) and the infusion of Aβ peptide (1–40) into the rat cerebral ventricle (right) on the number of RMEs evaluated by the radial maze task (see Materials and Methods for details). Left: Control rats (5% gum-arabic-administered rats; $n=21$) and EPA rats ($n=24$). After completing the initial behavior test, each of the two groups (control and EPA groups) was subdivided into two groups and infused with either (right) Aβ (Aβ group; $n=9$) or Aβ solvent (control group; $n=12$): the EPA group was divided into a vehicle-infused EPA group (EPA group; $n=12$) and an Aβ-infused EPA group (EPA+Aβ group; $n=12$). The four groups of rats were again behaviorally tested (with six trials) after the implantation of the miniosmotic pump. Each value represents the number of RMEs, presented as mean±S.E.M. in each block of six trials. The statistical significance of differences between the groups was determined by randomized two-factor (Block and Group) ANOVA followed by Bonferroni post hoc test. Groups without a common letter are significantly different at $P<0.05$. The mole percentages of the unsaturated fatty acids times the number of double bonds in each fatty acid are presented.

3.2. Effect of EPA on radial maze learning ability

250 The effect of the chronic administration of EPA on
251 working-memory- and reference-memory-related learning
252 ability is presented as the mean number of WMEs and RMEs
253 for each group, with data averaged over blocks of six trials
254 [Figs. 2 (left) and 3 (left), respectively]. Randomized two-
255 factor (Block and Group) ANOVA revealed a significant
256 main effect of both blocks of trials ($P=.0005$) and groups
257 ($P<.0001$), with a significant Block×Group interaction
258 ($P=.0018$), on the number of WMEs (Fig. 2, left). Similarly,
259 ANOVA revealed a significant main effect of both blocks of
260 trials ($P<.0001$) and groups ($P<.0001$), with a significant
261 Block×Group interaction ($P=.0484$), on the number of
262 RMEs (Fig. 3, left). These results indicate that EPA
263 administration improves working-memory- and reference-
264 memory-related learning ability in young rats.

265 The effect of EPA preadministered to the vehicle and Aβ-
266 infused groups on working-memory- and reference-memory-
267 related learning ability is presented as the mean number of
268 WMEs and RMEs for each group, with data averaged over
269 six trials [Figs. 2 (right) and 3 (right), respectively]. The
270 number of WMEs was significantly higher in the Aβ group
271 ($P=.0011$) than in the vehicle group (Fig. 2, right),
272 suggesting learning impairment — a well-known character-
273 istic of AD. The number of WMEs and RMEs was
274 significantly lower in the EPA+Aβ group (WMEs:
275 $P<.0001$; RMEs: $P=.0451$) than in the Aβ group [Figs. 2
276 (right) and 3 (right), respectively], indicating that preadmi-
277 nistration of EPA prevents cognitive deficits caused by the
278 infusion of Aβ into the cerebral ventricle of rats.

Table 1

Major fatty acid composition of plasma, cerebral cortex and hippocampus in control, EPA, Aβ and EPA+Aβ rats

	Control	EPA	Aβ	EPA+Aβ	
Plasma (%)					t1.3
EPA	0.36±0.01 ^c	4.53±0.29 ^a	0.48±0.04 ^c	3.80±0.21 ^b	t1.4
AA	26.7±0.86 ^a	21.5±0.68 ^b	25.2±0.46 ^a	22.0±1.17 ^b	t1.5
DHA	2.47±0.06 ^b	3.53±0.09 ^a	2.46±0.07 ^b	3.50±0.14 ^a	t1.6
DHA/AA	0.09±0.00 ^b	0.17±0.01 ^a	0.10±0.00 ^b	0.16±0.01 ^a	t1.7
USI	185±2.35 ^b	199±1.64 ^a	184±1.34 ^b	198±2.69 ^a	t1.8
Cortex (%)					t1.9
AA	12.0±0.23 ^{a,c}	11.1±0.11 ^b	12.4±0.41 ^a	11.4±0.18 ^{b,c}	t1.10
EPA	0.10±0.00 ^b	0.13±0.01 ^a	0.09±0.00 ^b	0.13±0.01 ^a	t1.11
DHA	16.8±0.29 ^{b,c}	18.1±0.17 ^a	16.1±0.44 ^b	17.6±0.29 ^{a,c}	t1.12
DHA/AA	1.41±0.05 ^b	1.63±0.03 ^a	1.32±0.08 ^b	1.56±0.05 ^a	t1.13
USI	167±1.15 ^b	173±0.86 ^a	164±1.48 ^b	171±1.20 ^a	t1.14
Hippocampus (%)					t1.15
AA	13.4±0.21 ^a	12.5±0.17 ^b	13.3±0.11 ^a	12.7±0.16 ^b	t1.16
EPA	0.10±0.00 ^b	0.14±0.01 ^a	0.10±0.00 ^b	0.14±0.00 ^a	t1.17
DHA	14.5±0.12 ^c	15.6±0.16 ^a	15.0±0.13 ^b	15.8±0.11 ^a	t1.18
DHA/AA	1.09±0.01 ^c	1.25±0.02 ^a	1.13±0.01 ^b	1.25±0.02 ^a	t1.19
USI	161±0.91 ^c	166±0.97 ^a	163±0.65 ^b	167±0.69 ^a	t1.20

Values are expressed as mean±S.E.M. and as mole percent of the total fatty acids ($n=9-12$). Means in a row with superscripts without a common letter differ ($P<0.05$). USI was calculated as a function of the sum of the mole percentages of unsaturated fatty acids times the number of double bonds in each fatty acid.

t2.1 Table 2

t2.2 Correlation between the mole percentages of plasma EPA and corticohippocampal EPA, DHA and USI

EPA (y)	Plasma x					
	Cortex			Hippocampus		
	EPA	DHA	USI	EPA	DHA	USI
t2.5	0.81	0.57	0.62	0.80	0.67	0.60
t2.6	(<i>P</i> <0.05)	(<i>P</i> <0.05)	(<i>P</i> <0.05)	(<i>P</i> <0.05)	(<i>P</i> <0.05)	(<i>P</i> <0.05)

t2.7 Results are evaluated with simple regression analysis. *P* values are expressed inside the parentheses.

279 3.3. Fatty acid profiles of plasma and brain

280 The major plasma fatty acid composition in the rat
 281 plasma, cortex and hippocampus is shown in Table 1. In the
 282 plasma, the proportion of EPA was significantly higher —
 283 and that of arachidonic acid [AA; 20:4(*n*-6)] was signifi-
 284 cantly lower (*P*<0.05) — in both EPA and EPA+Aβ rats than
 285 in the vehicle and Aβ rats, respectively. The proportion of
 286 DHA was higher in both EPA and EPA+Aβ rats than in the
 287 vehicle rats. EPA administration brought about a significant
 288 increase in the plasma DHA/AA molar ratio and USI value in
 289 both EPA and EPA+Aβ rats.

290 Chronic administration of EPA significantly enhanced
 291 the EPA proportion in both the cortex and the hippocampus
 292 of the EPA and EPA+Aβ rats. In the hippocampus, the
 293 proportion of DHA increased, whereas that of AA
 294 decreased significantly, effecting a significant increase in

the DHA/AA ratio in both the hippocampus and the
 cortex. EPA administration brought about a significant
 increase in the corticohippocampal USI values in both
 these groups of rats.

Highly significant positive correlations were observed
 between the percent compositions of plasma EPA and EPA,
 DHA or USI values of both the cortex and the hippocampus
 (Table 2), indicating that dietary administration of EPA
 accumulates EPA and DHA in brain tissues.

Table 3

Effects of chronic administration of EPA on hippocampal gene expression

Up-regulated genes	
1. Signal transduction	t3.4
Cystatin C	2.80 t3.5
GABA _B receptor1	2.04 t3.6
mGluR8	1.97 t3.7
Pyruvate dehydrogenase receptor 8	1.84 t3.8
Regulator of G-protein signaling 4	1.79 t3.9
2. Cell communication	t3.10
Syntaxin 1a	1.79 t3.11
PLD1	1.65 t3.12
Suppressor of K ⁺ transport defect 3	1.73 t3.13
Nucleic acid binding	t3.14
Forkhead box E1 (thyroid transcription factor 2)	2.38 t3.15
Basis helix–loop–helix domain containing class B2	2.02 t3.16
v-maf musculoaponeurotic fibrosarcoma (avian) oncogene family, protein G	1.72 t3.17
3. Microtubular dynamics	t3.18
VAPs B and C	1.67 t3.19
4. Enzyme	t3.20
VCP	2.07 t3.21
PLD1	1.65 t3.22
Protein arginine <i>N</i> -methyltransferase 3-like 3	1.60 t3.23
Pyridoxine 5-phosphate oxidase	1.59 t3.24
	t3.25
Down-regulated genes	t3.26
1. Signal transduction	t3.27
Insulin-like growth factor binding protein 2	2.09 t3.28
ATP-binding cassette, subfamily G (WHITE), member 5 (sterolin 1)	1.69 t3.29
GABA-α receptor γ3 subunit	1.63 t3.30
Hyperpolarization-activated cyclic nucleotide-gate K ⁺ channel 2	1.61 t3.31
2. Cell communication	t3.32
TTR	3.10 t3.33
Integrin α1	2.66 t3.34
Aquaporin 1	1.70 t3.35
Solute carrier family 9, member 1	1.63 t3.36
Transferrin receptor	1.61 t3.37
3. Nucleic acid binding	t3.38
Telomeric repeat binding factor 2	2.17 t3.39
ETS domain transcription factor Pet-1	1.80 t3.40
Myogenin	1.73 t3.41
Gonadotropin-inducible ovarian transcription factor 2	1.56 t3.42
4. Membrane component	t3.43
Ectonucleotide pyrophosphatase/phosphodiesterase 2	2.01 t3.44
5. Enzyme	t3.45
Coagulation factor 5	2.83 t3.46
Glycerol kinase	2.00 t3.47
Triadin 1	1.82 t3.48
Metalloprotease/disintegrin	1.78 t3.49
HP33	1.77 t3.50

Results are the average of triplicate determination, expressed as the ratio of EPA-administered rats to control rats. t3.51




Fig. 4. The effects of EPA on the LPO (upper; A and B) and ROS (lower; A and B) levels of the cortex (A) and hippocampus (B) of the Aβ rats. The correlation between corticohippocampal DHA/AA molar ratio and LPO levels (C) is presented.

330 The levels of both LPO and ROS were significantly
337 higher in the cerebral cortex of the A β rats (Fig. 4A) than in
338 the cerebral cortex of the vehicle controls and EPA rats. In
339 the cortex of the EPA+A β rats, both LPO and ROS levels
340 decreased, but decreased significantly only in the latter. In
341 the hippocampus of the A β rats, the level of LPO was
342 significantly higher than that in vehicle rats, but not
343 significantly higher than that in EPA rats (Fig. 4B). The
344 LPO levels were significantly lower in EPA+A β rats than in
345 A β rats. The ROS level in the hippocampus was also
346 significantly high in A β rats and significantly low in
347 EPA+A β rats. Finally, the change in oxidative stress in
348 these two tissues demonstrated a significant negative
349 correlation between LPO levels and corticohippocampal
350 DHA/AA ratios (Fig. 4C).

351 3.4. Effects of EPA administration on hippocampal 352 gene expression

353 EPA administration triggered a substantial change in the
354 expression of both up-regulated and down-regulated genes
355 (Table 3). The up-regulated genes strongly affected by EPA
356 were signal transduction proteins (cystatin C and GABA_B
357 receptor 1), cell communication protein (syntaxin 1a),
358 nucleic acid binding protein (thyroid transcription factor 2)
359 and microtubular dynamics [vesicle-associated membrane
360 proteins (VAPs) B and C]. The down-regulated genes
361 strongly affected by EPA were signal transduction protein
362 (ATP-binding cassette protein), cell communication proteins
363 [transferrin (TTR) and integrin α 1], nucleic acid binding
364 protein (telomeric repeat binding factor 2) and membrane-
365 associated ectonucleotidase.

366 4. Discussion

367 The present study provides evidence that not only the oral
368 administration DHA but also the oral administration of EPA
369 protects against memory impairment in the AD model rats
370 infused with amyloid peptide. The protective effect was
371 accompanied by corticohippocampal increases in EPA and
372 DHA, DHA/AA molar ratio and USI values. The 10- to
373 12-fold increase in the proportion of plasma EPA in the EPA
374 and EPA+A β rats (Table 1), as compared with that in the
375 controls, clearly suggests effective intestinal absorption of
376 this PUFA after oral administration. Moreover, the correla-
377 tion between plasma EPA and corticohippocampal DHA or
378 the DHA/AA ratio was highly positive, suggesting that
379 plasma EPA effectively deposits DHA in brain tissues after
380 crossing the blood–brain barrier.

381 We speculate that EPA increases the DHA (and DHA/AA
382 ratio) of the corticohippocampal tissues and, in doing so,
383 exerts beneficial effects on memory formation/protection in
384 EPA or EPA+A β rats. The basis of the speculation is that the
385 level of radiolabeled EPA ([¹⁴C]EPA) detected in the rat
386 brain 1 h after its oral administration to rats decreases time
387 dependently with a concomitant increase in the levels of

[¹⁴C]docosapentaenoic acid and [¹⁴C]DHA, metabolites of 388
EPA [13]. It is also inferred that each PUFA is metabolized 389
after being taken into cerebral endothelial cells and 390
astrocytes (constituent cells of the blood–brain barrier), 391
that it is released from both cells and that DHA is taken into 392
neurons as metabolite. This speculation is consistent with the 393
fact that DHA is taken into neurons from the extracellular 394
medium after its release from glial cells or the capillary 395
endothelium [14]. Indeed, increased DHA and EPA levels in 396
the plasma raise the fatty acid unsaturation index in rat 397
caudal arteries [18,19]. Moreover, lysophosphatidylcholine 398
may be a transporter of DHA in the blood–brain barrier [20]. 399
Remaining to be explored, however, is whether the 400
transformation of [¹⁴C]DHA from [¹⁴C]EPA occurs in the 401
liver and is then transported to the brain, or whether the 402
conversion occurs solely and/or partly in brain tissues. 403
Regardless of where EPA is transformed into DHA, it is 404
evident from our study that EPA was converted into DHA 405
and exerted its effects on memory functions. 406

The brain has an intrinsic capacity to retain its DHA; 407
however, if DHA is depleted, huge differences in brain 408
functions occur [21]. Thus, a change in brain DHA level 409
might be related to behavioral impairments [22]. We 410
reported that a small increase (in mol%) in DHA content 411
contributed significantly to limiting memory deficits in 412
DHA-deficient rats [8]. Thus, a small but highly significant 413
increase in corticohippocampal DHA composition (8– 414
9 mol%) in the EPA/EPA+A β rats after EPA administration 415
is consistent with our previous reports [8]. EPA-mediated 416
actions directly or indirectly could involve effects on 417
antioxidative status, amyloid processing, apoptosis, expres- 418
sion of a host of proteins, membrane lipid (disorder) 419
fluidity and exocytosis. An increased DHA/AA ratio is 420
associated with increased memory-related learning ability 421
in young [5], aged normal [6] and AD model rats [7,8], 422
with a concurrent decrease in brain LPO levels. Consistent 423
with our previous investigations [7,8], the DHA/AA ratios 424
correlated negatively with the repression of lipid peroxida- 425
tion (Fig. 4C). The mechanism by which this correlation 426
affects memory enhancement and amyloid burden is not 427
clear yet. Free-radical theory of AD pathology involves 428
amyloid-induced oxidative stress [23]. Increasing levels of 429
DHA in the cortex of aged rats significantly increase 430
antioxidative enzymes, including catalase, glutathione 431
peroxidase and reduced glutathione [24]. We have 432
hypothesized that the DHA/AA ratio acts as an indicator 433
of antioxidants indirectly by inhibiting the level of AA in 434
the neuronal plasma membrane [7]. An increase in the 435
DHA/AA ratio thus, at least partially, protects the 436
corticohippocampal regions from oxidative insult and 437
provides protection against the impairment of memory 438
in A β -infused rats. DHA inhibits the accretion of A β 439
peptide (1–40) in detergent insoluble neuronal membrane 440
domains of the cerebral cortex [25] and of A β -induced 441
apoptosis-like neuronal cell death [7]. Thus, the finding of 442
EPA-administration-induced protection against memory 443

444 impairment, with concurrent DHA accretion in the brain, is
445 in line with our studies [7,8] and those of others [10,26].

446 An alternative mechanism of EPA-induced amelioration
447 of memory may be as follows: we have previously reported
448 that DHA increases the expression of the Fos protein, the
449 immediate early gene *c-fos* (which acts as a transcription
450 factor and as a functional marker of neuronal activity) of the
451 rat CA1 hippocampus [27]. *n*-3 PUFAs induce the
452 expression of a host of genes that control synaptic plasticity
453 and underlying signal transduction mechanism(s) [11].
454 Using microarray analysis, we also examined whether the
455 administration of EPA induces the expression of genes, and
456 whether their function is correlated with EPA-induced
457 memory protection in the AD model rats. The gene chip
458 data are correlative and appear to be the only data from EPA-
459 versus-control rats in the present experimental scenario.
460 However, if such data had been obtained also from A β and
461 EPA+A β rats, then gene expression would have been
462 conferred to the neuroprotective actions of EPA in the
463 EPA+A β rats. Consistent with this possibility, we found that
464 EPA up-regulated 16 genes and down-regulated 25 genes
465 (Table 3). The γ -aminobutyric acid (GABA) receptor 1
466 (GABA_B) and the metabotropic glutamate receptor (mGluR)
467 1 were expressed twice, as compared with those of control
468 rats. The GABA_B receptor interacts with mGluR1-mediated
469 excitatory transmission [28], suggesting that EPA adminis-
470 tration might contribute to a mechanism of regulatory
471 synaptic plasticity. Several other genes that participate in
472 cell communication were also overexpressed. For example,
473 the expression of syntaxin 1a and VAPs B and C increased
474 concurrently (Table 3). Syntaxin 1a forms a complex with
475 VAP [29] in order to release neurotransmitters and regulates
476 their transporter [30]. These activities further support the
477 implication of the *n*-3 PUFA-induced increase in neuro-
478 transmitter release and synaptic plasticity. In addition, the
479 expression of the *cystatin C* gene, a lysosomal cysteine
480 protease inhibitor [31], increases upon EPA administration.
481 Lysosomal proteases (including cathepsins B and D) are up-
482 regulated in the AD brain [32], and cystatin C relates to
483 neurogenesis in neural stem cells [33]. In this regard, we
484 have recently shown that DHA significantly enhances
485 neurogenesis both in vivo and in vitro [34]. Phospholipase
486 D1 (PLD1) regulates both the release of secretory vesicles
487 from the trans-Golgi network (TGN) [35] and exocytosis
488 [36]. Amyloid proteolysis and subsequent trafficking from
489 the TGN to the cell surface are impaired in AD. Up-
490 regulation of PLD1 in AD rescues impaired APP trafficking
491 from TGN to the membrane surface [37]. In our study,
492 expression of the valosin-containing protein (VCP)
493 increased concomitantly in EPA-fed rats. VCP senses and
494 pulls abnormal protein accumulation in the cell, and pulls
495 from the endoplasmic reticulum and nucleus [38]. DHA-
496 induced clearance of A β from neuronal membranes is
497 suggested to be mediated by exocytosis — a process that
498 may be facilitated by the DHA-induced increase in
499 membrane fluidity [24,39,40]. Consistently, EPA-induced

memory augmentation is attributed to neurotransmitter 500
(noradrenaline) release [10], as facilitated by an increase in 501
membrane disorder (fluidity) [41]. A DHA-induced increase 502
in memory-related performance is accompanied by increased 503
levels of acetylcholine [42]. Therefore, all these effects are 504
speculated to be in concert with the EPA-induced increase in 505
memory protection of the EPA+A β rats. Our microarray data 506
also suggest that EPA suppressed the genes involved in 507
signal transduction, cell communication and/or membrane- 508
bound components to a remarkable extent (Table 3). The 509
contributions of EPA to the repression of these genes have 510
remained largely unrecognized. For example, TTR is a 511
thyroid hormone transporter that is secreted by the liver in 512
plasma and by choroid epithelial plexus in cerebrospinal 513
fluid. The TTR gene was suppressed in our feeding 514
paradigm. To date, the effects of *n*-3 PUFAs on the 515
expression of TTR are inconsistent and conflicting. Short- 516
term administration of *n*-3 fatty acids from fish oil for 517
1 month increases TTR expression in 2-year-old rats [43]. In 518
contrast, TTR gene expression is unchanged in mice fed a 519
high-DHA diet [44]. Our result is, however, consistent with 520
the report of Tanabe et al. [27] wherein the rats were fed *n*-3 521
PUFA containing 8% fish oil diet. In our study, the young 522
rats were fed 99% purified EPA at a dose of 300 mg/kg body 523
weight for 12 weeks. The discrepancies may thus relate to 524
the age of the rats, the feeding duration, the fatty acid 525
composition of the supplemented oil and the chow diet itself. 526
Also, the debate on whether TTR is expressed in the 527
hippocampus and/or whether the level found in brain tissues 528
is just an effect of contamination from the plexus epithelium 529
must be resolved [45]. Integrin α 1 expression decreased 530
consistently on EPA feeding. Inhibition of this cell- 531
communicating protein by echistatin or antibodies protects 532
against A β -induced degeneration of neurons in vitro [46]. 533
Because the gene chip data are unsubstantiated, with no 534
validation at either the mRNA level (by RT-PCR) or the 535
protein level, the correlation of EPA-induced memory 536
amelioration with other altered genes has remained without 537
further predication. The induction of gene expression thus 538
needs to be confirmed by real-time RT-PCR. 539

To summarize, EPA protects against A β -peptide-induced 540
memory deficit in AD model rats after its transformation into 541
DHA. This is accompanied by the accumulation of DHA 542
and/or an increase in the DHA/AA ratio in the corticohippo- 543
campal tissues, with a corresponding decrease in oxidative 544
stress and an increase in the expression of synaptic- 545
plasticity-related proteins. Nonetheless, further studies are 546
needed for additional data on EPA. 547

References 548

- [1] Hamano H, Nabekura J, Nishikawa M, Ogawa T. Docosahexaenoic 549
acid reduces GABA response in substantia nigra neuron of rat. 550
J Neurophysiology 1996;75:1264–70. 551
- [2] Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: 552
progress and problems on the road to therapeutics. *Science* 2002;297: 553
353–6. 554

- 555 [3] Kalmijn S, Launer LJ, Ott A, Witteman JC, Hofman A, Breteler MM. Dietary fat intake and the risk of incident dementia in the Rotterdam
556 Study. *Ann Neurol* 1997;42:776–82.
- 557 [4] Morris MC, Evans DA, Bienias JL, Tangney CC, Bennett DA, Wilson
558 RS, et al. Consumption of fish and *n*-3 fatty acids and risk of incident
559 Alzheimer disease. *Arch Neurol* 2003;60:940–6.
- 560 [5] Gamoh S, Hashimoto M, Sugioka K, Shahdat Hossain M, Hata N,
561 Misawa Y, et al. Chronic administration of docosahexaenoic acid
562 improves reference memory-related learning ability in young rats.
563 *Neuroscience* 1999;93:237–41.
- 564 [6] Gamoh S, Hashimoto M, Hossain S, Masumura S. Chronic adminis-
565 tration of docosahexaenoic acid improves the performance of radial arm
566 maze task in aged rats. *Clin Exp Pharmacol Physiol* 2001;28:266–70.
- 567 [7] Hashimoto M, Hossain S, Shimada T, Sugioka K, Yamasaki H, Fujii Y,
568 et al. Docosahexaenoic acid provides protection from impairment of
569 learning ability in Alzheimer's disease model rats. *J Neurochem* 2002;
570 81:1084–91.
- 571 [8] Hashimoto M, Tanabe Y, Fujii Y, Kikuta T, Shibata H, Shido O.
572 Chronic administration of docosahexaenoic acid ameliorates the
573 impairment of spatial cognition learning ability in amyloid beta-infused
574 rats. *J Nutr* 2005;135:549–55.
- 575 [9] Calon F, Lim GP, Yang F, et al. Docosahexaenoic acid protects from
576 dendritic pathology in an Alzheimer's disease mouse model. *Neuron*
577 2004;43:633–5.
- 578 [10] Song C, Horrobin D. Omega-3 fatty acid ethyl-eicosapentaenoate, but
579 not soybean oil, attenuates memory impairment induced by central IL-
580 1beta administration. *J Lipid Res* 2004;45:1112–21.
- 581 [11] Kitajka K, Puskas LG, Zvara A, Hackler Jr L, Barcelo-Coblijn G, Yeo
582 YK, et al. The role of *n*-3 polyunsaturated fatty acids in brain:
583 modulation of rat brain gene expression by dietary *n*-3 fatty acids. *Proc*
584 *Natl Acad Sci U S A* 2002;99:2619–24.
- 585 [12] Lauritzen L, Hansen HS, Jorgensen MH, Michaelson KF. The
586 essentiality of long chain *n*-3 fatty acids in relation to development
587 and function of the brain and retina. *Prog Lipid Res* 2001;40:1–94.
- 588 [13] Ishiguro J, Tada T, Ogiwara T, Murakami K, Kunihiro Y. Studies on the
589 metabolic disposition of ethyl eicosapentaenoate (EPA-E) in rats and
590 dogs. *Drug Metabol Dispos* 1987;2:683–702.
- 591 [14] Moore SA. Polyunsaturated fatty acid synthesis and release by brain-
592 derived cells in vitro. *J Mol Neurosci* 2001;16:195–200 [discussion
593 215–1].
- 594 [15] Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal
595 tissues by thiobarbituric acid reaction. *Anal Biochem* 1979;95:351–8.
- 596 [16] Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measure-
597 ment with the folin reagent. *J Biol Chem* 1951;193:265–75.
- 598 [17] Lockhart DJ, Dong H, Byrne MC, et al. Expression monitoring by
599 hybridization to high-density oligonucleotide arrays. *Nat Biotechnol*
600 1996;14:1675–80.
- 601 [18] Hashimoto M, Shinozuka K, Hossain MS, Kwon YM, Tanabe Y,
602 Kunitomo M, et al. Antihypertensive effect of all-*cis*-5,8,11,14,17-
603 icosapentaenoate of aged rats is associated with an increase in the
604 release of ATP from the caudal artery. *J Vasc Res* 1998;35:55–62.
- 605 [19] Hashimoto M, Shinozuka K, Gamoh S, et al. The hypotensive effect of
606 docosahexaenoic acid is associated with the enhanced release of ATP
607 from the caudal artery of aged rats. *J Nutr* 1999;129:70–6.
- 608 [20] Spector AA. Plasma free fatty acid and lipoproteins as sources of
609 polyunsaturated fatty acid for the brain. *J Mol Neurosci* 2001;16:
610 159–65 [discussion 215–1].
- 611 [21] Salem Jr N. Omega-3 fatty acids: molecular and biochemical aspects.
612 In: Spiller GA, Scala J, editors. *New protective roles for selected*
613 *nutrients*. New York: Alan R. Liss; 1989. p. 109–228.
- 614 [22] Salem Jr N, Moriguchi T, Greiner RS, McBride K, Ahmad A, Catalan
615 JN, et al. Alterations in brain function after loss of docosahexaenoate
616 due to dietary restriction of *n*-3 fatty acids. *J Mol Neurosci* 2001;16:
617 299–307.
- 618 [23] Behl C, Moosmann B. Antioxidant neuroprotection in Alzheimer's
619 disease as preventive and therapeutic approach. *Free Radic Biol Med*
620 2002;33:182–91.
- 621 [24] Hossain MS, Hashimoto M, Gamoh S, Masumura S. Antioxidative
622 effects of docosahexaenoic acid in the cerebrum versus cerebellum and
623 brainstem of aged hypercholesterolemic rats. *Neurochem* 1999;72:
624 1133–8.
- 625 [25] Hashimoto M, Hossain S, Agdul H, Shido O. Docosahexaenoic acid-
626 induced amelioration on impairment of memory learning in amyloid
627 beta-infused rats relates to the decreases of amyloid beta and
628 cholesterol levels in detergent-insoluble membrane fractions. *Biochim*
629 *Biophys Acta* 2005;1738:91–8.
- 630 [26] Lim S, Suzuki H. Changes in maze behavior of mice occur after
631 sufficient accumulation of docosahexaenoic acid in brain. *J Nutr* 2001;
632 131:319–24.
- 633 [27] Tanabe Y, Hashimoto M, Sugioka K, Maruyama M, Fujii Y, Hagiwara
634 R, et al. Improvement of spatial cognition with dietary docosahex-
635 aenoic acid is associated with an increase in Fos expression in rat CA1
636 hippocampus. *Clin Exp Pharmacol Physiol* 2004;131:700–3.
- 637 [28] Hirono M, Yoshioka T, Konishi S. GABA(B) receptor activation
638 enhances mGluR-mediated responses at cerebellar excitatory synapses.
639 *Neuroscience* 2001;12:1207–16.
- 640 [29] Fujiwara T, Yamamori T, Akagawa K. Suppression of transmitter
641 release by Tat HPC-1/syntaxin 1A fusion protein. *Biochim Biophys*
642 *Acta* 2001;1539:225–32.
- 643 [30] Sung U, Apparsundaram S, Galli A, Kahlig KM, Savchenko V,
644 Schroeter S, et al. A regulated interaction of syntaxin 1A with the
645 antidepressant-sensitive norepinephrine transporter establishes cate-
646 cholamine clearance capacity. *J Neurosci* 2003;23:1697–9.
- 647 [31] Abrahamson M, Barrett AJ, Salvesen G, Grubb A. Isolation of six
648 cysteine proteinase inhibitors from human urine. *J Biol Chem* 1986;
649 261:11282–9.
- 650 [32] Cataldo AM, Barnett JL, Pieroni C, Nixon RA. Increased neuronal
651 endocytosis and protease delivery to early endosomes in sporadic
652 Alzheimer's disease: neuropathologic evidence for a mechanism of
653 increased β -amyloidogenesis. *J Neurosci* 1997;17:6142–51.
- 654 [33] Taupin P, Ray J, Fischer VM, Suhr ST, Hakansson K, Grubb A, et al.
655 FGF-2-responsive neural stem cell proliferation requires CCg, a novel
656 autocrine/paracrine cofactor. *Neuron* 2000;28:385–7.
- 657 [34] Kawakita E, Hashimoto M, Shido O. Docosahexaenoic acid promotes
658 neurogenesis in vitro and in vivo. *Neuroscience* 2006;139:991–7.
- 659 [35] Chen YG, Siddhanta A, Austin CD, Hammond SM, Sung TC,
660 Frohman MA, et al. Phospholipase D stimulates release of nascent
661 secretory vesicles from the trans-Golgi network. *Cell Biol* 1997;138:
662 495–504.
- 663 [36] Humeau Y, Vitale N, Chasserot-Golaz S, Dupont JL, Du G, Frohman
664 MA, et al. A role for phospholipase D1 in neurotransmitter release.
665 *Proc Natl Acad Sci U S A* 2001;98:15300–5.
- 666 [37] Cai D, Zhong M, Wang R, Netzer WJ, Shields D, Zheng H, et al.
667 Phospholipase D1 corrects impaired beta APP trafficking and
668 neurite outgrowth in familial Alzheimer's disease-linked presenilin-
669 1 mutant neurons. *J Cell Death Differ* 2001;8:977–84.
- 670 [38] Hirabayashi M, Inoue K, Tanaka K, Nakadate K, Ohsawa Y, Kamei Y,
671 et al. *Cell Death Differ* 2001;8:977–84.
- 672 [39] Shahdat H, Hashimoto M, Shimada T, Shido O. Synaptic plasma
673 membrane-bound acetylcholinesterase activity is not affected by
674 docosahexaenoic acid-induced decrease in membrane order. *Life*
675 *Sci* 2004;74:3009–24. ~~*Proc Natl Acad Sci U S A* 2006;103:
676 1936–0.~~
- 677 [40] Hashimoto M, Hossain S, Shimada T, Shido O. Docosahexaenoic acid-
678 induced protective effect against impaired learning in amyloid beta-
679 infused rats is associated with increased synaptosomal membrane
680 fluidity. *Clin Exp Pharmacol Physiol* 2006;33:934–9.
- 681 [41] Clarke MS, Prendergast MA, Terry Jr AV. Plasma membrane ordering
682 agent pluronic F-68 (PF-68) reduces neurotransmitter uptake and
683 release and produces learning and memory deficits in rats. *Learn Mem*
684 1999;6:634–9.
- 685 [42] Minami M, Kimura S, Endo T, Hamaue N, Hirafuji M, Togashi H, et al.
686 Dietary docosahexaenoic acid increases cerebral acetylcholine levels
687 and improves passive avoidance performance in stroke-prone
688

Q5

Q6

Q7

- 689 spontaneously hypertensive rats. *Pharmacol Biochem Behavior* 1997; 690 58:1123–9.
- 691 [43] Puskás LG, Kitajka K, Nyakas C, Barcelo-Coblijn G, Farkas T. Short- 692 term administration of omega 3 fatty acids from fish oil results in 693 increased transthyretin transcription in old rat hippocampus. *Proc Natl 694 Acad Sci U S A* 2003;100:1580–5.
- 695 [44] Lim GP, Calon F, Morihara T, Yang F, Teter B, Ubeda O, et al. A diet 696 enriched with the omega-3 fatty acid docosahexaenoic acid reduces amyloid burden in an aged Alzheimer mouse model. *J Neurosci* 2005; 697 25:3032–40. 698
- [45] Sousa JC, Cardoso I, Marques F, Saraiva MJ, Palha JA. Transthyretin 699 and Alzheimer's disease: where in the brain? *Neurobiol Aging* 2007; 700 28:713–8. 701
- [46] Anderson KL, Ferreira A. Integrin activation A link between β - 702 amyloid deposition and neuronal death in aging hippocampal neurons. 703 *J Neurosci Res* 2004;75:688–97. 704 705

706

UNCORRECTED PROOF