

Seasonal Influenza Infection Risk Factors of Personnel at Shimane University Hospital

Yumi JO¹⁾, Masahiro HIROSE²⁾, Toshihiko KAWAMURA³⁾, Chiaki SANO⁴⁾, Shunichi KUMAKURA⁵⁾, Junji SUZUMIYA⁶⁾, Kiwamu NAGOSHI⁷⁾

¹⁾ Infection Control Department, Shimane University Hospital, Izumo, Shimane 693-8501, Japan

²⁾ Department of Community-based Health Policy and Quality Management, Faculty of Medicine, Shimane University, Izumo, Shimane 693-8501, Japan

³⁾ Department of Medical Informatics, Shimane University Hospital, Izumo, Shimane 693-8501, Japan

⁴⁾ Department of Community Medicine Management, Faculty of Medicine, Shimane University, Izumo, Shimane 693-8501, Japan

⁵⁾ Department of Medical Education & Research, Faculty of Medicine, Shimane University, Izumo, Shimane 693-8501, Japan

⁶⁾ Innovative Cancer Center, Shimane University Hospital, Izumo, Shimane 693-8501, Japan

* Present Institution: Sunkohkai-Kouga Hospital, 2-30-1 Daikakuji, Yaizu, Shizuoka 425-0088, Japan

⁷⁾ Department of Environmental Medicine and Public Health, Faculty of Medicine, Shimane University, Izumo, Shimane 693-8501, Japan

(Received April 9, 2021; Accepted April 26, 2021)

Objectives: A questionnaire survey is performed to investigate the factors associated with seasonal influenza infection at Shimane university hospital. **Methods:** From 2014 to 2017, questionnaire were distributed to hospital employees who were eligible for vaccination. The sheet includes the following information: age of responders, living situation with children, their pre-season vaccination status, and their history of seasonal influenza infection. **Results:** We analyzed data of a total of 5450 employees, and 6.2% had seasonal influenza infection history in the preceding season. Female sex (odds ratio [OR] 1.364, $P < 0.027$), aged 30-39 years (vs 20-29 years, OR 2.023, $P < 0.001$; vs ≥ 60 years; OR 4.206, $P < 0.001$), and living with children (OR 2.191, $P < 0.001$) were identified as factors associated with risk of influenza infection. **Conclusion:** Recognition of these risk factors may be beneficial to prevent and control influenza infection, and hospital employees should pay utmost attention for avoiding secondary influenza infection from households.

Keywords: seasonal influenza infection; hospital employee; influenza vaccination; household infection

INTRODUCTION

Seasonal influenza infection control in hospitals is important for securing patient safety. An annual influenza vaccination for hospital employees has the potential to benefit their patients and their families as well as hospital administrators. Hospital employees often come into contact with patients, children, and older adults. The Annual vaccination is recommended for hospital employees to avoid contracting seasonal influenza and spreading it to others, especially hospitalized patients. According to the WHO fact sheet, the “seasonal flu vaccine is the most effective method of preventing infection.”

Influenza vaccination has been reported to have protective efficacy for healthy adults even if the vaccine differs from the subtype of the prevalent virus. Furthermore, influenza vaccination decreases the risk of complications and death among elderly people, children, and patients with underlying diseases, despite inadequate prevention [1-5]. In Japan, the vaccinations are performed once or twice a year for each applicant.

According to the website of the Ministry of

Corresponding author: Yumi Jo, M.D.

Infection Control Department, Faculty of Medicine, Shimane University, 89-1 Enya-Cho, Izumo, Shimane 693-8501 Japan

Tel: +81-853-20-2483

Fax: +81-853-20-2428

E-mail address: joyumi@med.shimane-u.ac.jp

Health, Labor and Welfare (MHLW), influenza vaccination does not entirely prevent influenza infection but has a potent effect on the aggravation of the disease. However, the reported decrease in the incidence of influenza infection due to vaccination varies among studies [6-10]. Recently, an increasing number of immunocompromised patients and the spread of new influenza virus strains have led to an upsurge in the vaccination rate. Some reports have indicated that high-dose vaccination or twice inoculation can be useful for influenza prevention [11, 12]. It is difficult to provide the high-volume inoculation and twice inoculation for the hospital personnel in the season when vaccine is short in the market.

Moreover, some hospital employees have to be absent because of influenza infection despite undergoing vaccination, and this absence has a bad influence on their regular duties. In our hospital, prophylactic medication is prescribed at the hospital's expense for hospital employees who come into frequent contact with influenza patients without taking infection precautions. However, prophylactic medication is not common in Japan.

At our hospital which has already conducted vaccination and prophylactic medication use, some hospital employee has influenza. The aim of this observational study was to investigate factors associated with influenza infection and measures that can reduce the impact of influenza on hospital personnel.

METHODS

Study Design

A questionnaire survey was conducted to hospital employees at Shimane university hospital, Japan, between the last four seasonal influenza vaccination campaigns (2014/2015, 2015/2016, 2016/2017, 2017/2018). Shimane University Hospital is a ter-

tiary, referral and advanced technology hospital with 600 beds approved by the MHLW in Japan.

Hospital employees who participated in the seasonal influenza vaccination campaign were enrolled in the study and received a questionnaire sheet. Seasonal influenza is prevalent from December until April of the following year. In the hospital, the seasonal influenza vaccination campaign starts on October 30 and ends by the second week in November. A questionnaire survey has been administered since 2014 for the purpose of investigating the factors associated with an influenza infection situation. The questionnaire sheet was distributed within two days before the date of seasonal influenza vaccination and collected with a preliminary examination sheet on the day of inoculation. This questionnaire included the next items; 1) age, 2) the gender, 3) occupation, 4) whether the person had been vaccinated in the preceding fiscal year (preseason vaccination status), 5) whether the individual lived with children aged less than 15 years, 6) the respondent's history of seasonal influenza infection in the preceding season. We excluded the responders who did not answer the question about the history of influenza infection. For diagnosing influenza infection, employees in whom diagnosis was not made by positive results of rapid antigen test were excluded from "infected" and were categorized as "unsure".

Statistical Analyses

JMP14.2® software (SAS Institute Japan) and STATA14.0® software (STATA Corp USA) were used for statistical analysis. A chi-squared test was used for the comparison between the two groups. Odds ratio for the influenza infection risk was calculated, and predictive factors for influenza infection were determined by using logistic regression analysis. The level of significance was set at $p < 0.05$.

Table 1. Number of vaccinated subjects, number of vaccines used and questionnaire responses
Valid response, the number of responses that was excluded the responders who did not answer the question about the history of influenza infection.

Year	2014	2015	2016	2017	Total
Vaccine applicants	1676	1700	1853	1901	7130
Responders to questionnaires	1111	1455	1642	1683	5891
Eligible responders*	1091	1408	1452	1499	5450

*Exclude the responders who did not answer the question about the history of influenza

RESULTS

Basic Characteristics

The number of vaccine inoculators and responders to questionnaire during study periods is shown in Table 1. Of 5891 responders, we excluded 441 responders who did not answer the question about the history of influenza infection, data of a total of 5450 employees were analyzed.

The characteristics of the responders in this study are shown in Table 2. Of 5450 responders, 2194 were nurses, 724 were physicians, 868 were office workers, and 1580 were other employees (including laboratory technicians and other medical staff). 5052 received influenza vaccine and 366 did not receive in the preceding fiscal year. 1716 lived with children, and 2829 did not live with children.

Table 2. Characteristics of questionnaire responders
With children, respondents living with children aged <15 years old; without children, respondents not living with children aged <15 years old; vaccinated, vaccinated responders; unvaccinated, nonvaccinated responders

	Number
Gender*	
Male	1267
Female	4099
Age (years old)**	
20-29	1554
30-39	1489
40-49	1106
50-59	909
≥ 60	341
Occupation†	
Nurse	2194
Physician	724
Office worker	868
Others	1580
Vaccination§	
Vaccinated	5052
Un-vaccinated	366
Living situation¶	
With children	1716
Without children	2829
Influenza infection‡	
Infected	333
Uninfected	5053
Unsure	64

Excludes unknown number as below; 84 (*), 51 (**), 84 (†), 32 (§), 905 (¶)

For diagnosing influenza infection, employees in whom diagnosis was not made by positive results of rapid antigen test were excluded from "infected" and were categorized as "unsure".

Incidence Rate of Seasonal Influenza Infection

The seasonal influenza infection incidence during the whole period was 6.2%. The incidence rate was

5.3% in 2013/2014, 4.9% in 2014/2015, 6.7% in 2015/2016, and 7.5% in 2016/2017.

The incidence rates of the seasonal influenza infection were significantly different among sexes (Table 3). The total infection rate of females for all 4 seasons was significantly higher ($p = 0.027$, Odds Ratio (OR) 1.364, 95% Confidence Interval (95% CI) 1.027-1.812) (Table 3).

A significant difference in the distribution of infected subjects was found according to age (Table 3). The incidence rate of seasonal influenza infection was the highest among subjects in their 30s in all seasons.

As we show in Table 4, level 1 and level 2 logistic regression analysis was performed among age generations, and the corresponding odds ratios were determined.

Employees aged 30-39 years were more susceptible to seasonal influenza infection as compared with those aged 20-29 years (OR 2.023, 95% CI 1.509-2.712, $P < 0.001$), and those aged ≥ 60 years (OR 4.206, 95% CI 2.040-8.669, $P < 0.001$) (Table 4).

No difference was found in the incidence according to occupation.

Vaccination Situation

The incidence rate of seasonal influenza infection was 6.2% in vaccinated employees and 5.8% in unvaccinated employees ($p = 0.75$) (Table 3). The annual infection incidence rates were 5.3% in the vaccinated employees and 5.4% in the unvaccinated employees ($p = 0.99$) in 2013/2014, 4.9% in the vaccinated employees and 4.9% in the unvaccinated employees ($p = 1.00$) in 2014/2015, 6.8% in the vaccinated employees and 5.1% in the unvaccinated employees ($p = 0.57$) in 2015/2016, and 7.5% in the vaccinated employees and 8.7% in the unvaccinated employees ($p = 0.67$) in 2016/2017, and no significant differences were observed between the groups.

Living Situation with/without Children Younger than 15 Years Old

The incidence rate of seasonal influenza was higher in respondents living with children aged < 15 years old (With-children) than in those not living with children aged < 15 years old (Without-children) (p

Table 3. Number and percentage of individuals with seasonal influenza among hospital staff by vaccination status, gender, living situation with children, and age, relative risk and odds ratio Shimane, Japan, September 1, 2013-August 20, 2017

Infected means people who contracted seasonal influenza infection.

Uninfected means people who did not contract seasonal influenza infection.

Vaccinated, vaccinated respondents; unvaccinated, nonvaccinated respondents; with children, respondents living with children aged <15 years old; without children, respondents not living with children aged <15 years old; OR: Odds ratio; 95% CI: 95% confidence interval.

	Infected no.(%)	Uninfected no.(%)	<i>P</i> -value	OR (95%CI)
Gender				
Male	62 (4.9)	1193 (95.1)		1
Female	268 (6.6)	3780 (93.4)	0.0274	1.364 (1.027-1.812)
Age (years old)				
20-29	74 (4.8)	1473 (95.2)	< 0.0001	
30-39	135 (9.2)	1328 (90.8)		
40-49	67 (6.2)	1017 (93.8)		
50-59	45 (5.0)	857 (95.0)		
≥ 60	8 (2.4)	331 (97.6)		
Vaccination				
Vaccinated	311 (6.2)	4683 (93.8)	0.7527	
Un-vaccinated	21 (5.8)	340 (94.2)		
Living situation				
With children	154 (9.2)	1529 (90.9)	< 0.0001	2.191 (1.715-2.800)
Without children	123 (4.4)	2677 (95.6)		

OR: Odds ratio; 95% CI: 95% confidence interval

Table 4. Level 1 and level 2 logistic regression analysis of age generation and corresponding odds ratios

Level 1	Level 2	<i>P</i> -value	OR	95%CI
20-29	30-39	< 0.0001	2.023	1.509-2.712
20-29	40-49	0.1182	1.311	0.933-1.842
20-29	50-59	0.8196	1.045	0.714-1.528
20-29	≥ 60	0.0524	0.481	0.229-1.007
30-39	20-29	< 0.0001	0.494	0.368-0.662
30-39	40-49	0.0052	0.648	0.478-0.878
30-39	50-59	0.0002	0.516	0.364-0.731
30-39	≥ 60	< 0.0001	0.237	0.115-0.490
40-49	20-29	0.1182	0.762	0.542-1.071
40-49	30-39	0.0052	1.543	1.138-2.091
40-49	50-59	0.2525	0.797	0.540-1.175
40-49	≥ 60	0.0082	0.366	0.174-0.771
50-59	20-29	0.8196	0.956	0.654-1.398
50-59	30-39	0.0002	1.935	1.366-2.742
50-59	40-49	0.2525	1.254	0.850-1.850
50-59	≥ 60	0.0462	0.46	0.214-0.986
≥ 60	20-29	0.0524	2.078	0.992-4.352
≥ 60	30-39	< 0.0001	4.206	2.040-8.669
≥ 60	40-49	0.0082	2.725	1.295-5.733
≥ 60	50-59	0.0462	2.172	1.013-4.657

OR: Odds ratio; 95% CI: 95% confidence interval

< 0.001) (Table 3). During 2015-2017, the infection rates were markedly higher in the With-children group than in the Without-children group.

The annual infection incidence rates were 7.0% in

the With-children group and 4.5% in the Without-children group in 2013/2014, with no significant difference ($p = 0.09$); 7.8% in the With-children group and 3.3% in the Without-children group ($p <$

0.001, OR 2.46, 95%CI 1.44-4.23) in 2014/2015; 10.5% in the With-children group and 4.4% in the Without-children group ($p < 0.001$, OR 2.55, 95%CI 1.52-4.33) in 2015/2016; and 11.1% in the With-children group and 5.7% in the Without-children group ($p = 0.001$, OR 2.11, 95%CI 1.32-3.39) in 2016/2017.

Relationship between Vaccination Situation and Living-with-Children Situation

In the With-children group, the infection rates during four seasons in the vaccinated and unvaccinated were 9.1% and 8.9%, respectively, and there was no significant difference between groups ($p = 0.95$).

Simultaneously, in the With-children group, the annual infection incidence rates in the vaccinated and unvaccinated groups were 7.3% and 2.8% in 2013/2014, 7.6% and 11.1% in 2014/2015, 10.3% and 13.0% in 2015/2016, and 11.2% and 11.5% in 2016/2017, respectively, and no significant differences were observed between the two groups (p values: 0.31 in 2013/2014, 0.50 in 2014/2015, 0.68 in 2015/2016, and 0.96 in 2016/2017).

In the Without-children group, the infection rates in the vaccinated and unvaccinated groups were 4.4% and 4.2%, respectively, with no significant difference between groups ($p = 0.86$).

Moreover, the annual incidence rates in the vaccinated and unvaccinated groups were 4.5% and 5.8% ($p = 0.61$) in 2013/2014, 3.4% and 1.9% ($p = 0.56$) in 2014/2015, 4.8% and 0.0% in 2015/2016, and 5.5% and 8.2% ($p = 0.44$) in 2016/2017, respectively, and there were no significant differences between the two groups.

DISCUSSION

The Relationship between Influenza Infection and Living-with-Children Situation

This report determined the relationship between seasonal influenza infection and living-with-children situation by using statistical method.

On the basis of the chi-squared quadratic profile, there were significant differences in the incidence of influenza infection by age (Table 3). A significant difference was found for hospital employees in their 30s (Table 4). Next to 30s, the person in the 40s

had influenza at a higher percentage as compared with the person of other generations. When we added an investigation, the living together rate with children was found to be 66% in the 30s group. The living together rate with children was 57% in the 40s group, and the rate was 15% in the 20s group, and the rate was 15% in the 50s group, and the rate was 10% in the 60s group. It may influence that prevalence of influenza is high in 30s group that a rate to live together with children is high. It is inferred that the difference in prevalence by the generation is due to the difference in living together rate with children.

The incidence rate of seasonal influenza infection for females was significantly higher than that for males (6.62% for females and 4.94% for males; $p = 0.03$), and the odds ratio of influenza infection for females was 1.364 (95% CI: 1.03-1.81). A living together rate with the children of males was 37.1%, and the rate of females was 37.9%. It cannot be said that the frequency that females live together with children is higher than males ($P = 0.33$). According to the hospital employees' comments on the questionnaire sheets, many female hospital employees stated that they did not want to part from their children. According to some reports, 37-51.7% of families share a bed until a child reaches approximately 12 years old, which is the age of a primary schoolchild in Japan [13-17]. We cannot judge it from this study, but the behavior of female staff in the home may be different from it of male staff. The people living together with children seem to increase clearly at an opportunity to come in contact with children.

The association between the influenza infection of children and the influenza infection of the hospital personnel was not able to evaluate in this investigation. Why is the person living together with children easy to have influenza? The reason may be that children living together had influenza and may be the difference of the lifestyle (e.g., the participation of the children's association) by there being children. In future, we want to investigate more why a person living together with children tend to get influenza.

Influenza infection risk of the vaccinated respond-

ers

The incidence rates of female were significantly high. The incidence rates were 5.0% for males and 6.7% for females ($p = 0.042$, OR 1.36, 95%CI 1.01-1.86). The incidence rate was the highest among subjects in their 30s, with rates of 9.3%. A significant difference in the distribution of infected subjects was found according to age ($p < 0.001$). The annual infection incidence rates were 9.1% in the With-children group and 4.4% in the Without-children group. The incidence rates of the With-children group were significantly high ($p < 0.001$, OR 2.16, 95%CI 1.66-2.81)

It is important for hospital employees to reduce opportunities against influenza virus exposure both at home and in the workplace. In other words, to control hospital-acquired infection, hospital employees need to be educated about influenza infection prevention in outside of the hospital.

According to the report, the prophylactic administration of oseltamivir (75 mg/day for 4 days) could prevent hospital employees from developing influenza after contact with patients with influenza [18]. If prophylaxis is recommended when the staff came into contact with patients with influenza in home, the influenza infection of the hospital personnel may decrease. In addition to influenza vaccination, precautions of wearing a mask and hand hygiene should be promoted during the influenza season [19]. Separating and isolating family members with influenza from healthy family members at home may be an effective measure to prevent influenza infection. It should be further considered whether vaccination of children could reduce the incidence of seasonal influenza in hospital personnel.

The incidence of influenza infection is reduced by employee influenza vaccination, and vaccination reduces sick leave of employee [20]. From our results, the next phenomenon seems to a fact. We cannot prevent influenza infection by a certain one measure, for example a "vaccination". The hospital personnel should perform many precautions against influenza infection in parallel not to get seasonal influenza. In 2020/2021, mask wearing and hand washing, securing of sociological distance are recommended to not only the hospital personnel but also a citizen for prevention of infection of the new

coronavirus infection. We are getting a custom to avoid a crowd as a new lifestyle. We could not know the change of the real vaccination rate, but vaccination was strongly recommended. An infected person begins to increase the seasonal flu from December in an average year, but the number of the reports of influenza does not show a tendency of the increase at the end of January in 2020/2021 season.

Limitations

This study has several limitations. Some factors associated with influenza infection differ during the four seasons.

Firstly, it is generally thought that the efficacy of vaccines might be different according to season. We could not evaluate differences in influenza vaccine efficacy between hospital personnel and individuals who do not work in the hospital. Secondly, we were not able to accurately determine the number of subjects present when the questionnaire was administered. The subject of a distributor and the respondent of the questionnaire became extended. Third, precautions taken against infection vary among families. Therefore, it is difficult to conclude that all people living with children are at increased risk of influenza infection.

Finally, frozen section diagnosis was not performed in all individuals with fever. A prospective study is necessary to evaluate the effect of the vaccination.

CONCLUSIONS

For hospital employees living with children younger than 15 years old, the odds ratio of seasonal influenza infection was 2.19. They were at increased risk of the influenza infection. It is suggested that it reduce an incidence of the seasonal influenza infection among another employee and patients at a hospital that a staff takes the infection-related precautionary measures at home. In other words, influenza vaccination is necessary for the prevention of influenza infection, but even a person inoculated vaccine into has influenza. It is also important to reduce influenza virus exposure opportunity anytime. Next is inferred by an influenza outbreak trend in 2020/2021.

Social behavior modification and the rise in vaccination rate reduce the influenza infection of the hospital personnel.

Ethics Statement

This study was approved by the Medical Ethics Committee at Shimane University Faculty of Medicine “Shimane University Institutional Committee on Ethics” on October 10, 2018. The ethics code is 20180914-1.

Contributors

Conceived of and designed the experiments: YJ, CS, and SK. Performed the experiments: YJ, TK and MH. Analyzed the data: YJ and MH. Wrote the paper: YJ, MH, JS and SK. Managed the study: KN

Acknowledgments

The authors thank all of the respondents who completed the questionnaire and Sakane K. and Ishitobi E. for help with the vaccinations.

Funders

None.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

REFERENCES

- 1) Wilhelm M. Influenza in older patients: A call to action and recent updates for vaccinations. *Am. J. Manag. Care* 2018;24:S15–S24.
- 2) Jefferson T, Rivetti A, Harnden A, Pietranonj CD, Demicheli V. Vaccines for preventing influenza in healthy children. *Cochrane Database Syst Rev* 2008;2: CD004879. doi:10.1002/14651858.CD004879.pub3.
- 3) Santos GD, Tahrat H, Bekkat-Berkani R. Immunogenicity, safety, and effectiveness of seasonal influenza vaccination in patients with diabetes mellitus: A systematic review. *Hum Vaccin Immunother* 2018;14:1853–66. doi:10.1080/21645515.2018.1446719.
- 4) Bosaeed M, Kumar D. Seasonal influenza vaccine in immunocompromised persons. *Hum Vaccin Immunother* 2018;14:1311–22. doi:10.1080/21645515.2018.1445446.
- 5) Bitterman R, Eliakim-Raz N, Vinograd I, Trestioreanu AZ, Leibovici L, Paul M. Influenza vaccines in immunosuppressed adults with cancer. *Cochrane Database Syst Rev* 2018;2:CD008983. doi:10.1002/14651858.CD008983.pub3.
- 6) Rondy M, Kissling E, Emborg HD, *et al.* Interim 2017/18 influenza seasonal vaccine effectiveness: Combined results from five European studies. *Euro Surveill* 2018;23:18–00086. doi:10.2807/1560-7917.ES.2018.23.9.18-00086.
- 7) Flannery B, Chung JR, Belongia EA, *et al.* Interim estimates of 2017-18 seasonal influenza vaccine effectiveness - United States, February 2018. *MMWR Morb Mortal Wkly Rep* 2018;67:180–5. doi:10.15585/mmwr.mm6706a2.
- 8) Chiu SS, Kwan MYW, Feng S, *et al.* Interim estimate of influenza vaccine effectiveness in hospitalised children, Hong Kong, 2017/18. *Euro Surveill* 2018;23: 18–00062. doi:10.2807/1560-7917.ES.2018.23.8.18-00062.
- 9) Mira-Iglesias A, López-Labrador FX, Guglieri-López B, *et al.* Influenza vaccine effectiveness in preventing hospitalisation of individuals 60 years of age and over with laboratory-confirmed influenza, Valencia Region, Spain, influenza season 2016/17. *Euro Surveill* 2018;23:17–00318. doi:10.2807/1560-7917.ES.2018.23.8.17-00318.
- 10) Ishikane M, Kamiya H, Kawabata K, *et al.* Seasonal influenza vaccine (A/New York/39/2012) effectiveness against influenza A virus of health care workers in a long term care facility attached with the hospital, Japan, 2014/15: A cohort study. *J Infect Chemother* 2016;22:777–9. doi: 10.1016/j.jiac.2016.03.011.
- 11) Young-Xu Y, Aalst RV, Mahmud SM, *et al.* Relative vaccine effectiveness of high-dose versus standard-dose influenza vaccines among veterans health administration patients. *J Infect Dis* 2018;217: 1718–27. doi:10.1093/infdis/jiy088.
- 12) Cordero E, Roca-Oporto C, Bulnes-Ramos A, *et al.* Two doses of inactivated influenza vaccine improve immune response in solid organ transplant recipients: Results of TRANSGRIPE 1–2, a randomized controlled clinical trial. *Clin Infect Dis* 2017;64:829–38. doi:10.1093/cid/ciw855.

- 13) Yamada F, Fujiwara T. Prevalence of self-reported shaking and smothering and their associations with co-sleeping among 4-month-old infants in Japan. *Int J Environ Res Public Health* 2014;11:6485–93. doi:10.3390/ijerph110606485.
- 14) Asaka Y, Takada S. Comparing sleep measures of infants derived from parental reports in sleep diaries and acceleration sensors. *Acta Paediatr* 2011;100:1158–63. doi:10.1111/j.1651-2227.2011.02204.x.
- 15) Nelson EAS, Taylor BJ, Jenik A, *et al.* International child care practices study: Infant sleeping environment. *Early Hum Dev* 2001;62:43–55. doi:10.1016/s0378-3782(01)00116-5.
- 16) Mindell JA, Sadeh A, Wiegand B, How TH, Goh DYT. Cross-cultural differences in infant and toddler sleep. *Sleep Med* 2010;11:274–80. doi:10.1016/j.sleep.2009.04.012.
- 17) Fukumizu M, Kaga M, Kohyama J, Hayes MJ. Sleep-related nighttime crying (yonaki) in Japan: A community-based study. *Pediatrics* 2005;115:217–24. doi:10.1542/peds.2004-0815c.
- 18) Nagao S, Onishi M, Nishiyama C. Study of the use of oseltamivir in the prevention and treatment of 2009/H1N1 influenza. *Jpn J Infect Prev Control* 2012;27:178–82. doi:10.4058/jsei.27.178. (Eng Abstr)
- 19) Aiello AE, Murray GF, Perez V, *et al.* Mask use, hand hygiene, and seasonal influenza-like illness among young adults: A randomized intervention trial. *J Infect Dis* 2010; 201:491–8. doi:10.1086/650396.
- 20) Susa R, Fuse K, Ishizawa M, Nakamata M, Tsukada H, Gejyo F. Effectiveness of influenza vaccine in hospital workers. *Jpn J Infect Prev Control* 2001;16:303–8. doi:10.11550/jsei1986.16.303. (in Japanese)