

A Doppler Flow Analysis of the Neovascular Flow in Carotid Plaque Using Ultrasound Micro-flow Imaging

Fumio NAKAGAWA, Hidemasa NAGAI, Shoko URUSHIMATSU, Yuta FUJIWARA,

Mizuki KAMBARA, Tsutomu YOSHIKANE, Takeshi MIYAZAKI, Yasuhiko AKIYAMA

Department of Neurosurgery, Faculty of Medicine, Shimane University, Izumo, Shimane 693-8501, Japan

(Received December 24, 2020; Accepted January 15, 2021)

Carotid artery stenting (CAS) was recommended for the treatment of severe carotid artery stenosis. One problem associated with CAS is cerebral infarction due to embolic events. Vulnerable plaque with bleeding and neovascularization is a risk factor for such events. Bleeding in plaque has been evaluated by ultrasonography and MRI. However, detection of neovascularization in plaque has not been established. We measured the intraplaque microcirculatory blood flow of 6 patients who underwent CAS with superb micro-vascular imaging (SMI). We compared the SMI signal and the embolic events following CAS. Case 3 had an SMI-positive signal with clear Doppler flow in the affected plaque; he subsequently showed plaque protrusion and asymptomatic embolic events. Another patient showed no Doppler signal. Doppler flow detected by SMI was found to be related to neovascularity of plaque. This finding might indicate the plaque vulnerability.

Keywords: superb micro-vascular imaging, carotid artery stenosis, neovascularization, carotid artery stenting

INTRODUCTION

Internal carotid artery stenosis accounts for approximately 20% of all cases of cerebral infarction; the major cause of stenosis is atheromatous plaque. The current treatments for carotid artery stenosis include medical treatment and surgical treatment. If carotid artery stenosis is severe or symptomatic, carotid artery stenting (CAS) or carotid endarterectomy (CEA) is recommended as an evidence-based treatment for carotid stenosis.

Embolic events occur as perioperative complications when performing CAS. Emboli are caused by debris generated from vulnerable plaque. Debris may be generated in CAS procedures, such as the passing a guidewire or device through a stenotic location, stent placement, and/or balloon dilation. Alternatively, the debris may occur when plaque is pushed out of the mesh of the stent; this is called plaque protrusion.

Preoperative vulnerable plaque is currently attracting attention as a risk factor for embolic complications. Plaque vulnerability is associated with intraplaque hemorrhage and neovascularization within the plaque. We have already evaluated plaque bleeding by carotid ultrasound and magnetic resonance (MR) carotid plaque imaging. However, no method of evaluation has been established for the examination of neovascularization in plaque. In order to examine neovascularity, contrast-enhanced ultrasound (CEUS) studies have been conducted; however, the use of contrast agents in Japan poses problems, such as lack of insurance coverage for carotid artery imaging, and possible side effects associated with the use of contrast agents [1, 2]. On the other hand, superb micro-vascular imaging (SMI) is a recent ultrasound technology that can display images comparable to CEUS [3, 4].

Corresponding author: Fumio Nakagawa, MD, PhD

Department of Neurosurgery, Faculty of Medicine, Shimane University, 89-1 Enya-cho, Izumo, Shimane 693-8501, Japan

E-mail: fumio@med.shimane-u.ac.jp

Tel: +81-853-20-2245

Fax: +81-853-21-8954

SMI is a new blood flow imaging technique with high sensitivity, high resolution, and low artifact [5, 6]. The technique can effectively separate blood flow signals and overlapping tissue motion artifacts by the adaptive calculation method, and accurately detect low-speed blood flow signals [7].

We are investigating the possibility of assessing and quantifying plaque vulnerability, which is a risk factor for CAS, by capturing the microblood flow of neovascularization around the plaque and measuring Doppler signaling by carotid ultrasonography without contrast medium.

MATERIALS AND METHODS

Patients

This was a single-center, prospective observational study. The subjects included 6 consecutive patients who underwent CAS for internal carotid artery stenosis at Shimane University from January 2018 to April 2018. Patients who could not undergo carotid ultrasound, CAS, or filter protection were excluded.

The CAS procedure was performed using a modified Parodi method by special neuro-endovascular doctors (FN and YA). A 9-Fr guiding catheter with a balloon was placed in the common carotid artery and the external carotid artery was occluded by a Percutaneous Occlusion Balloon™ (Medtronic, Minneapolis, MN, USA), and a Filter Wire EZ™ (Boston Scientific, Natick, MA, USA) was used as a filter protection device. Precise ProRX™ (Cordis, Miami, FL, USA) was used as the stent. The diameter of stent was a target blood vessel diameter of +1 to 2 mm and the stent was designed for stenosis of $\geq +2$ cm in length.

The present study was improved by the institutional review board of the Shimane University Medical Ethics Committee (Management number: 20171114-1, approval number: 2950), and participants received a written informed consent.

Methods

An Aplio i900™ (Canon Medical Systems Corporation, Tochigi, Japan) ultrasound machine was used. The probe used was the same model as the wide-band linear probe PLI-705BX™. The ultrasound was performed as a preparative test for CAS. The fol-

lowing parameters were measured: peak systolic flow velocity (PSV), end-diastolic velocity (EDV), mean flow velocity (MFV), inner diameter of blood vessel, and maximum intima-media thickness (maxIMT) of the internal carotid artery. The stenosis rate was calculated using the ECST method and the area method. In addition to these measured values, ultrasound micro-flow imaging by SMI was performed. It was considered that a high echo signal on SMI indicated neovascularization in the plaque [1]. Then, if a clear Doppler signal was detected at the each high echo signal site of SMI, we captured a clear blood flow signal and defined it as a positive Doppler signal. In order to unify the ultrasound procedure, measurements were taken by the same neurosonographer (HN) and the results were confirmed by two doctors (FN and HN).

In addition, as a preoperative evaluation, ulcer formation was evaluated by measuring the diameter, length, and shape of the stenosis on 3D-CT angiography (CTA). Vulnerability was evaluated based on the presence or absence of intra-plaque bleeding by MR carotid plaque imaging and time-of-flight (TOF)-MR angiography. The angiographic findings and cone beam CT were used to evaluate plaque protrusion. Moreover, the occurrence of a region of high signal intensity was evaluated by postoperative MR-diffusion-weighted imaging (DWI).

We investigated and analysed the preoperative items (ultrasound, MRI, and CTA), the intraoperative findings (cerebral angiography), and the postoperative items (MR-DWI).

RESULTS

One of the six patients who participated in the study was excluded due to difficulty in stent placement. The remaining 5 cases included 4 males and 1 female, with an average age of 75.2 years; all patients had unilateral lesions (Table 1). Each item was measured for 10 blood vessels, with the blood vessels treated with CAS defined as the target blood vessels and the untreated blood vessels defined as controls (Table 2).

In the target blood vessels, the average flow velocity was 223.8 cm/s (42.3–337.3 cm/s) for PSV, 76.6 cm/s (19.0–165.6 cm/s) for EDV, and 86.9

Table 1. Examination data on the lesion side

Case	1	2	3	4	5	Average
Age	74	73	82	70	77	75.2
Gender	M	F	M	M	M	
Disease side	Left	Right	Left	Left	Left	
Blood vessel inner diameter (mm)	5.9	6.8	5.6	5.6	6.4	6.1
maxIMT (mm)	4.2	4.2	5.5	5.0	3.1	4.4
PSV (cm/s)	337.3	42.3	222.8	315.5	201.0	223.8
EDV (cm/s)	165.6	19.0	72.1	68.7	57.6	76.6
Vmean (cm/s)	-	27.5	125.5	-	107.7	86.9
ECST (%)	76	68	82	70	65	72.2
Area (%)	97	-	90	85	79	87.8
Echo brightness	Low	High	High	Low	Low	
SMI high echo signal	+	+	+	+	+	
SMI Doppler signal	-	-	+	-	-	
CTA ulceration	+	+	-	+	-	
MR TOF	ISO	LOW	ISO	HIGH	LOW	
plaque image	T1WI -	ISO	ISO-HIGH	ISO	ISO-HIGH	
	T2WI -	ISO	ISO-HIGH	HIGH	ISO-HIGH	
Plaque Protrusion	-	-	+	-	-	
Postoperative DWI	-	-	+	-	-	

Table 2. Examination data on the contralateral side

Case	1	2	3	4	5	Average
Contralateral side	Right	Left	Right	Right	Right	
Blood vessel inner diameter (mm)	5.2	6.6	5	4.6	5.4	5.4
maxIMT (mm)	2.6	2.8	5.2	1.7	1.8	2.8
PSV (cm/s)	81.6	67.7	84.2	129.3	84.8	89.5
EDV (cm/s)	22.9	31.2	35.7	39	26	31.0
Vmean (cm/s)	43.3	51.6	42.2	74.5	45.7	51.5
Echo brightness	HIGH	HIGH	HETEROGENOUS	LOW	HIGH	
SMI high echo signal	-	-	-	-	-	
SMI Doppler signal	-	-	-	-	-	

cm/s (27.5–125.5 cm/s) for MFV. The average stenosis rate was 72.2% (65–82%) in ECST and 87.75% (79–97%) in area. The average plaque thickness was 4.4 mm (3.1–5.5 mm) at max IMT. A positive of the high echo signal of SMI in the plaque was detected in all 5 cases. A clear Doppler signal at the SMI-positive site could not be detected in the plaque, with the exception of Case 3. A clear Doppler signal was detected in this case, which was only bidirectional during systole (Fig. 1).

In the control blood vessels, the average flow velocities were 89.5 cm/s (67.7–129.3 cm/s) for PSV, 31.0 cm/s (22.9–39 cm/s) for EDV, and 51.5 cm/s (42.2–74.5 cm/s) for MFV. The stenosis rate was not calculated for the control blood vessels. The average plaque thickness was 2.8 mm (1.7–5.2 mm) at max IMT. The high echo signal of SMI was negative in all 5 cases and there was no apparent hyperintensity in the plaque (Tables 1 and 2).

The preoperative examination of the target vessels revealed ulceration by CTA in 3 of 5 cases, and MR plaque imaging suggested bleeding in 2 of 5 cases.

Intraoperative plaque protrusion was only seen in 1 (Case 3) of 5 cases. Case 3, postoperative MRI revealed high intensity spots on DWI. On the other hand, preoperative CTA was revealed no ulceration, and MR plaque imaging showed some high signals in the plaque. Case 3, SMI showed positive high echo signals and positive Doppler signals in the plaque (Fig. 1). In this plaque protrusion case, the Doppler signal findings were observed at the three sites of SMI high echo signal (Figs. 1 and 2). In addition, for comparison with MR plaque imag-




Fig. 1. ① Doppler signal-positive findings and the detection site on SMI in Case 3





Fig. 2. ②③ Doppler signal-negative findings and the detection site on SMI in Case 3

ing, the corresponding positions of the three sites of SMI high echo signal were shown. (Figs. 1 and 2 and 3). However, there was no apparent relationship among the SMI high echo signal, the SMI Doppler signal, and the high intensity spot on MR-DWI.

DISCUSSION

Carotid artery stenting was recommended for severe carotid stenosis with atheromatous plaque. Carotid

plaques not only cause the stenosis and the occlusion of blood vessels, but also cause arterial embolism due to plaque vulnerability [8, 9]. Vulnerable plaque was one of the surgical indications for CAS and CEA, even in cases in which the stenosis ratio was not so severe. Conversely, vulnerable plaque is associated with a high risk of stroke during the stenting procedure. It is also known that vulnerable plaque is prone to complications when performing CAS, including plaque protrusion, distal embolus,




Fig. 3. Plaque imaging (T1 FS) in Case 3
 ① Shows the Doppler signal-positive part in Fig.1.
 ② and ③ Show the Doppler signal-negative parts in Fig.2.




Fig. 4. Doppler signal-negative findings and detection sites on SMI in cases other than Case 3

and clogging of the filter, which was a distal embolus protection device. For this reason, it is important to understand how to easily detect vulnerable plaque in preparation for CAS. The pathological findings of vulnerable plaque were characterized as intraplaque bleeding, necrotic tissue, and fibrous cap rupture. On CTA vulnerable plaque showed ulceration, which was a sign that represented plaque rupture. MR plaque imaging show signal intensity on T1 and T2, which represents bleeding. The reported ultrasound findings of carotid vulnerable plaque include low echo brightness, ulceration, and mobile plaque. [10, 11]. These findings from the evaluation of carotid plaque vulnerability by carotid ultrasound depended on the performance of the ultrasonic equipment, the frequency of the probe, and the professional skill of the sonographer.

Yamamoto *et al.* noted, "The SMI had made it possible to extract clinically necessary slow blood flow information by identifying features specific to motion artifacts and enhancing their ability to eliminate them. As a result, the boundary with the blood flow cavity could be made clearer than the conventional blood flow display method, and the morphology and shape could be evaluated in more detail"[12]. Therefore, the morphology of the carotid plaque could be evaluated more easily and in detail using SMI. In addition to the superiority in the evaluation of the plaque morphology, SMI has been attracting attention as a test that can visualize low-flow blood flow without using contrast medium [3, 4]. Zhang *et al.* and Mahtab *et al.* noted, "A significant proportion of ischemic strokes are caused by emboli from unstable carotid artery plaques with intra-plaque neovascularization as a key feature of plaque instability. When intra-plaque neovascularization was evaluated using SMI and CEUS and compared with pathological findings, there was a correlation between the density of intra-plaque neovascularization and the findings of SMI and CEUS" [1, 13]. Intra-plaque neovascularization was reported to be associated with the promotion of the inflammatory reaction, increased permeability of the fibrous cap, increase of the atheroma core, and intra-plaque hemorrhage [14]. These findings of neovascularization are suggested to be possible causes of plaque vulnerability.

In this study, routine imaging modalities such as CTA, MR plaque imaging and carotid ultrasonography did not reveal any specific results in any of the 5 cases. Thus, we did not think that all patients had a high risk of CAS before surgery. However, in Case 3, plaque protrusion occurred at the time of stent procedure, and asymptomatic embolic lesions were seen by DWI postoperatively. In this protrusion case, the preoperative findings included high echo plaque on carotid ultrasound, no ulceration on CTA, ISO intensity on MR-TOF, ISO to high intensity on T1WI of MR plaque imaging, and high echo signal on SMI. These findings were also observed in the other 4 cases. The present case differed from the other cases in the positive Doppler signal at the site of the plaque that showed a high echo signal on SMI. We hypothesized that the difficult conditions might have occurred for the following reasons. 1) The neovascularization could not be displayed directly by routine carotid ultrasound, CTA, MR-TOF or MR plaque imaging. 2) In this study we observed very low flow of the SMI high echo signal in the plaque in the affected vessel that was a candidate for CAS. It was reported that the SMI high echo signal was neovascularization [1]. Hence if it was suggested that the SMI high echo signal represented neovascularization, we thought that this was a relatively common condition in candidates for CAS. 3) However, if this signal included noise, the true neovascularization was indicated by a clear Doppler signal. Doppler signal assessment of intraplaque neovascularization by SMI provides information on plaque vulnerability that cannot be detected by other tests.

The present study was associated with several limitations. It was performed in a single center study and the study population was small, so it lacks statistical power. However, being able to more accurately grasp plaque vulnerability using ultrasonic equipment with SMI, which can be easily performed in routine medical care, plays an important role in deciding the treatment policy. Thus, we would like to continue to collect cases and show their statistical effectiveness.

In conclusion, plaque with a positive SMI Doppler signal might indicate true neovascularization, and it was considered that the plaque with the clear

SMI Doppler signal was highly vulnerable.

REFERENCES

- 1) Zhang H, Du J, Wang H, *et al.* Comparison of diagnostic values of ultrasound micro-flow imaging and contrast-enhanced ultrasound for neovascularization in carotid plaques. *Exp Ther Med* 2017;14:680-8. doi: 10.3892/etm.2017.4525.
- 2) Saito K, Nagatsuka K, Ishibashi-Ueda H, Watanabe A, Kannki H, Iihara K. Contrast-enhanced ultrasound for the evaluation of neovascularization in atherosclerotic carotid artery plaques. *Stroke* 2014;45:3073-5. doi: 10.1161/STROKEAHA.114.006483.
- 3) Song ZZ, Zhang YM. Contrast-enhanced ultrasound imaging of the vasa vasorum of carotid artery plaque. *World J Radiol* 2015;7:131-3. doi: 10.4329/wjr.v7.i6.131.
- 4) Burke AP, Farb A, Malcom GT, Liang YH, Smialek J, Virmani R. Coronary risk factors and plaque morphology in men with coronary disease who died suddenly. *N Engl J Med* 1997;336:1276-82. doi: 10.1056/NEJM199705013361802.
- 5) Staub D, Partovi S, Schinkel AFL, *et al.* Correlation of carotid artery atherosclerotic lesion echogenicity and severity at standard US with intraplaque neovascularization detected at contrast-enhanced US. *Radiology* 2011;258:618-26. doi: 10.1148/radiol.10101008.
- 6) Russell DA, Wijeyaratne SM, Gough MJ. Changes in carotid plaque echomorphology with time since a neurologic event. *J Vasc Surg* 2007;45:367-72. doi: 10.1016/j.jvs.2006.09.048.
- 7) Sztajzel R, Momjian S, Momjian-Mayor I, *et al.* Stratified gray-scale median analysis and color mapping of the carotid plaque: correlation with endarterectomy specimen histology of 28 patients. *Stroke* 2005;36:741-5. doi: 10.1161/01.STR.0000157599.10026.ad.
- 8) Yuan C, Zhang SX, Polissar NL, *et al.* Identification of fibrous cap rupture with magnetic resonance imaging is highly associated with recent transient ischemic attack or stroke. *Circulation* 2002;105:181-5. doi: 10.1161/hc0202.102121.
- 9) Murphy RE, Moody AR, Morgan PS, *et al.* Prevalence of complicated carotid atheroma as

- detected by magnetic resonance direct thrombus imaging in patients with suspected carotid artery stenosis and previous acute cerebral ischemia. *Circulation* 2003;107:3053-8. doi: 10.1161/01.CIR.0000074204.92443.37.
- 10) Yamagami H. B mode, intergrated backscatter and plaque properties. In: The Japan Academy of Neurosonology. *Manual of Neurosonology*. 1st ed. Tokyo: Hokosya; 2006:114-119.
- 11) Kume S, Hama S, Yamane K, Wada S, Nishida T, Kurisu K. Vulnerable carotid arterial plaque causing repeated ischemic stroke can be detected with B-mode ultrasonography as a mobile component: Jellyfish sign. *Neurosurg Rev* 2010;33:419-30. doi: 10.1007/s10143-010-0270-9.
- 12) Yamamoto T. Usefulness of SMI in the vascular area. *Eizojo Medical* 2018;50:45-52.
- 13) Zamani M, Skagen K, Scott H, Lindberg B, Russell D, Skjelland M. Carotid plaque neovascularization detected with superb microvascular imaging ultrasound without using contrast media. *Stroke* 2019;50:3121-7. doi: 10.1161/STROKEAHA.119.025496.
- 14) Doyle B, Caplice N. Plaque neovascularization and antiangiogenic therapy for artherosclerosis. *J Am Coll Cardiol* 2007;49:2073-80. doi: 10.1016/j.jacc.2007.01.089.