

# 学位論文の要旨

氏名 豊田 元哉

学位論文名 Association of Mild Kidney Dysfunction With Silent Brain Lesions in Neurologically Normal Subjects

発表雑誌名 Cerebrovascular Diseases EXTRA, in press  
(巻, 初頁～終頁, 年)

著者名 Genya Toyoda, Hirokazu Bokura, Shingo Mitaki, Keiichi Onoda, Hiroaki Oguro, Atsushi Nagai, Shuhei Yamaguchi

## 論文内容の要旨

### **INTRODUCTION**

Recently, the relation between chronic kidney disease (CKD) and cerebrovascular disease has been highlighted in not only symptomatic but also asymptomatic cases. Several population studies have reported that there is an independent association of estimated glomerular filtration rate (eGFR) with silent brain infarction (SBI), white matter lesions, and microbleeds (MBs), which are important independent factors for poor prognosis of patients with CKD. Although a large number of studies reported the relationship between each silent brain lesion and CKD, there are few reports in which comprehensive analyses were performed for the association of CKD with all types of subclinical brain lesions. Thus, the aim of this study was to investigate the relationship between all silent brain lesions and CKD, and provide new insights regarding the relative significance of CKD to each brain lesion in a large-scale, cross-sectional, neurologically normal population.

### **MATERIALS AND METHODS**

We studied a total of 1,937 subjects (1,012 men and 925 women) with a mean age of  $59.4 \pm 7.9$  years (range: 27 to 86 years). The screening system included collection of medical, neurological, and psychiatric history, formal neurological examinations by an experienced neurologist, neuropsychological assessment, MRI of the head, and blood and urine analyses. The inclusion criteria for this study were the following: no history of neurological or psychiatric disorders, no abnormalities on neurological examination, no severe medical illness, and written

informed consent to participate in this study. The study design was approved by the institutional ethics committee of Shimane University Hospital.

Venous blood samples were collected from all subjects after an overnight fast, and the sera were used for measurements of fasting blood glucose, HbA1c, lipid, and creatinine using an autoanalyzer. The level of urinary protein was examined and we defined positive proteinuria when the dipstick scale showed a value  $\geq 1+$ . Kidney function was estimated by the calculated creatinine clearance using the 4-variable Modification of Diet in Renal Disease equation. CKD was defined as either positive proteinuria or eGFR  $< 60$ . Subjects with an eGFR  $< 30$  were not included in the present study.

MRI scans of the brain were performed using a 1.5-Tesla MRI system (Symphony Ultra Gradient, Siemens). SBI was defined as a focally hyperintensity lesion  $> 3$  mm in diameter in the T2WI, corresponding to a hypointensity lesion in the T1WI. PVH was graded using a scale of 0 to 4, and we defined grade 3 and 4 as PVH +. SWML was graded using a scale of 0 to 3 according to the Fazekas' grading scale, and we defined grades 2, 3, and 4 as SWML +. MBs were defined as homogenous round foci of signal loss that were 2–10-mm in diameter in T2\*WI.

Baseline characteristics were compared between the CKD and non-CKD group using the student's *t*-test for parametric data and the Mann-Whitney U-test for non-parametric data. The relationship between MRI changes and CKD was analyzed using the Pearson's chi-squared test. *P* values adjusted for age and sex were also given for univariate analysis. After adjustment for age and sex, the multivariate logistic models were adopted to estimate the risks of the presence of CKD for silent brain lesions.

## **RESULTS AND DISCUSSION**

Of 1,937 subjects, the prevalence of CKD was 8.7%. The mean age of CKD subjects was significantly higher than the non-CKD subjects. After adjusted for age and sex, CKD was associated with hypertension and dyslipidemia. All silent lesions on MRI, including SBI, subcortical white matter lesion (SWML), periventricular hyperintensity (PVH), and MBs were more prevalent in subjects with CKD after adjusted for age and sex. Age had a strong effect on all brain lesions. After adjustment for age and sex, hypertension was associated with the presence of all brain lesions except for PVH, and diabetes mellitus also affected the presence of SBI and MBs, whereas dyslipidemia was not related to any silent brain lesions.

With binary logistic regression analysis, the presence of CKD was a significant risk factor for all types of silent brain lesions independent of other risk factors. The odd's ratio was the highest for MBs among the silent brain lesions. Although hypertension was the strongest risk factor for SBI, other brain lesions were affected more strongly by the presence of CKD compared to hypertension.

The current study demonstrated the independent and significant role of CKD on silent brain lesions in addition to traditional vascular risk factors. To our best knowledge, this is the largest

cross-sectional study that examined the relationship of CKD to structural brain changes in a general healthy population. Not only univariate analysis but also multivariate logistic regression analysis showed aggravating influence of CKD on any silent brain lesions independent of other confounding vascular risk factors.

The current study demonstrated that even mild CKD is associated with SBI independently of hypertension and age. Among all silent brain lesions only SBI was affected more strongly by hypertension than CKD (odd's ratio: 2.31 vs 1.90), and this pattern was also reported in a recent study. The relatively strong contribution of hypertension to SBI indicates that blood pressure control may be critically important to prevent SBI in subjects with mild CKD.

Although hypertension is the strongest risk factor for SBI, other MRI lesions, including PVH, SWML, and MBs, are more strongly influenced by the presence of CKD than hypertension in the current study. There were some differences in the relationship between CKD and SWML or PVH. PVH was affected only by age and CKD, whereas SWML was affected by hypertension in addition to age and CKD. This differential association might be attributable to the histopathological features of PVH, which involve the disruption of the ependymal lining with the sub-ependymal widening of the extracellular space.

MBs are considered clinically silent but are strongly associated with advanced small vessel or microvascular disease. Previous reports have examined the relationship between MBs and CKD in specific populations such as patients with intracranial hemorrhage, those with hypertension, or those with severe kidney dysfunction. The current study demonstrated that even mild kidney dysfunction could be a risk factor for the appearance of MBs independent of hypertension in a general population. Furthermore, MBs was affected by CKD most strongly among silent brain lesions. Since MBs are a strong risk factor for future cerebral hemorrhage and ischemic stroke, active interventions that prevent kidney dysfunction may be important for reducing occurrence of future stroke in subjects with risk factors for CKD.

## **CONCLUSION**

The current study demonstrated that even mild CKD is associated with all types of silent brain lesions independent of age, sex, hypertension, and diabetes mellitus in neurologically normal population. Specifically MBs was most strongly associated with the presence of CKD. Further prospective studies are required to define the causal relationship between silent brain lesions and CKD.

論文審査及び最終試験又は学力の確認の結果の要旨

甲・㉔	氏名	豊田 元哉
学位論文名	Association of Mild Kidney Dysfunction with Silent Brain Lesions in Neurologically Normal Subjects	
学位論文審査委員	主査 副査 副査	石橋 豊 廣田 秋彦 田邊 一明
<p>論文審査の結果の要旨</p> <p>近年、慢性腎臓病（CKD）が症候性のみならず無症候性脳血管障害と密接な関連があることが注目され、無症候性脳病変の存在はCKD症例における独立した予後規定因子であると報告されているが、CKDと無症候性脳病変の関連についての研究は、十分ではない。本研究では、CKDが無症候性脳病変の独立した危険因子であるか否かを大規模横断的に詳細に検討した。神経学的に健常で推算糸球体濾過量（estimated glomerular filtration rate: eGFR）30 ml/min/1.73m<sup>2</sup>以上の1,937名（男1,012名、女925名、平均年齢59歳）を対象に、CKDの有無と無症候性脳病変の関係を検討した。CKDは、尿蛋白1+以上かeGFR &lt; 60と定義した。脳病変は、頭部MRI画像から無症候性脳梗塞（silent cerebral infarction: SBI）、白質病変としてのperiventricular hyperintensity（PVH）とsubcortical white matter lesion（SWML）、微小出血（microbleeds: MBs）の有無を調べ、無症候性脳病変におけるCKDおよび年齢、性別（男性）、喫煙の有無、飲酒習慣の有無、教育歴（年数）、高血圧症の有無、糖尿病の有無、脂質異常症の有無を因子として解析を行った。その結果、喫煙、飲酒習慣、教育歴、脂質異常を除く因子において脳病変に対して有意な関与が認められた。CKDは対象の8.75%で認められ、CKD群において全ての脳病変が有意に多かった。CKD、年齢、性別、高血圧症、糖尿病を因子とした脳病変に対する多変量ロジスティック解析では、性別は有意ではなかったが、高血圧症はPVHを除く3つの病変と、糖尿病はSBIとMBsと、年齢およびCKDはすべての脳病変と有意な関連を認めた。CKDは、他の因子の影響を補正しても有意であり、特にMBs病変において最も強いオッズ比を示した（OR 3.30: 1.51-7.20, p&lt;0.005）。すなわち、CKDは、無症候性脳病変に対する独立した危険因子となり得ることが判明した。その機序として脳動脈と腎動脈に共通するstrain vesselの血管内皮傷害が関与することが考えられた。本研究は、CKDを有する例においては積極的な脳病変検索と予防管理の重要性を示唆しており、今後の更なる治療戦略への発展を期待させるものであり、博士（医学）の学位授与に値すると判断した。</p> <p>最終試験又は学力の確認の結果の要旨</p> <p>申請者は、脳ドックを受検した1,937名を対象としてMRIによる無症候性脳病変と慢性腎臓病の有無および各種動脈硬化因子との関連を検討し、慢性腎臓病が無症候性脳病変の独立した危険因子であることを示した。これまでの報告に比して対象人数は最も多く、得られた詳細な解析結果は、慢性腎臓病を有する例での積極的な予防管理の重要性を示す説得力のある論文である。関連分野の知識も十分に有しており、医学博士の学位授与に値すると判断した。（主査:石橋 豊）</p> <p>本研究は、無症候性脳病変と軽度な慢性腎臓病との関係を1,937名という大きい母集団で調べ、MRIにより診断されたSBI、PVH、SWML、MBsの4つの因子が慢性腎臓病を増悪させる独立した危険因子であることをestimated GFRを指標に用いて示したものである。近い将来の臨床応用見据えた研究内容は、臨床系の学位に相応しく、質疑応答からは基礎知識も十分あることが伺え、学位授与に値すると判断した。（副査 廣田秋彦）</p> <p>申請者は、脳ドックを受検した1937例を対象として、慢性腎臓病が無症候性脳病変の独立した危険因子であることを示した。高血圧を含む心臓血管危険因子の積極的な予防管理の重要性を示す臨床的に価値のある論文である。関連分野の知識も十分に有し、学位授与に値すると判断した。（副査 田邊一明）</p>		

（備考）要旨は、それぞれ400字程度とする。