

1 **A Mesenchymal Stem cell line (B10) increases angiogenesis in a rat MCAO**  
2 **model**

3

4 Abdullah Md. Sheikh<sup>a</sup>, Shozo Yano<sup>a</sup>, Shingo Mitaki<sup>b</sup>, Md. Ahsanul Haque<sup>b</sup>, Shuhei  
5 Yamaguchi<sup>b</sup>, Atsushi Nagai<sup>a</sup>.

6

7 <sup>a</sup>Department of Laboratory Medicine and <sup>b</sup>Department of Neurology, Shimane  
8 University School of Medicine, 89-1 Enya Cho, Izumo 693-8501, Japan.

9

10

11 Corresponding Author:

12 Atsushi Nagai,

13 Department of Laboratory Medicine

14 Shimane University School of Medicine

15 89-1 Enya Cho,

16 Izumo 693-8501

17 Japan

18 Tel: +81-0853-20-2312

19

20

1 **Abstract:**

2 A human mesenchymal stem cell line (B10) transplantation has been shown to  
3 improve ischemia-induced neurological deficits in animal stroke models. To  
4 understand the underlying mechanism, we have investigated the effects of B10  
5 transplantation on cerebral angiogenesis in a rat middle cerebral artery occlusion  
6 (MCAO) model. B10 cells were transplanted intravenously 24 h after MCAO.  
7 Immunofluorescence staining results showed that compared to PBS-group, vWF  
8 positive vessel and endoglin positive new vessels were increased in B10-  
9 transplanted MCAO groups in the lesion areas. The mRNA of angiogenesis factors  
10 including placental growth factor and hypoxia inducible factor (HIF)-1 $\alpha$  were  
11 increased 3 days after MCAO in the core and IBZ areas of B10-transplanted group.  
12 Angiopoetin1 mRNA was increased only in the IBZ. Western blotting results  
13 showed that HIF-1 $\alpha$  and vascular endothelial growth factor (VEGF) proteins levels  
14 were increased in B10-transplanted group. Double immunofluorescence staining  
15 results revealed that both HIF-1 $\alpha$  and VEGF were expressed in  
16 macrophage/microglia in the core area. In the IBZ, however, HIF-1 $\alpha$  was  
17 expressed both in astrocytes and macrophage/microglia, while VEGF was  
18 expressed only in macrophage/microglia. Moreover, TGF $\beta$  protein levels were  
19 found to be increased in B10-transplanted group in the core and IBZ regions. Cell  
20 culture experiments using a human microglia cell line (HMO6) and B10 showed  
21 that IL-1 $\beta$  induced VEGF mRNA expression in both cell types. IL-1 $\beta$  was found to  
22 be highly expressed in B10 cells, and its co-culture with HMO6 further increased

1 that in B10. Co-culture increased VEGF mRNA in both B10 and HMO6. In the rat  
2 brains, IL-1 $\beta$  was expressed in macrophage/microglia and transplanted-B10 cells  
3 in the core. IL-1 $\beta$  positive cell number was increased slightly, but significantly in  
4 B10-transplanted rats. To explore further, IL-1 $\beta$  expression was silenced in B10  
5 cells by transfecting mRNA specific siRNA, and then transplanted in MCAO rats.  
6 Immunostaining result showed that although total endoglin positive area was  
7 decreased, vessel-like structure appeared as early as 3 days after MCAO when IL-  
8 1 $\beta$ -silenced B10 was transplanted. Thus our results demonstrated that B10 cells  
9 increased angiogenesis in MCAO rat model, through the regulation of HIF-1 $\alpha$  and  
10 VEGF expression, where IL-1 $\beta$  might play a role.

11 Key words: Mesenchymal stem cell line (B10); transplantation; angiogenesis;  
12 MCAO; HIF-1 $\alpha$ ; VEGF; IL-1 $\beta$

13

## 1 **Introduction:**

2 Brain tissue is highly dependent on oxygen for its energy metabolism, and  
3 consequently can tolerate oxygen deprivation only for a short period of time  
4 (Bélanger et al., 2011; Kumar et al., 2010). Cerebral vasculature is developed  
5 uniquely by making an elaborate arteriolar anastomotic network. Such vascular  
6 arrangement helps to protect brain tissue during conditions of decreased blood  
7 supply through anastomotic collateral circulation (Liebeskind, 2003). Stroke occurs  
8 due to severe interruption of cerebral blood flow that generally results from  
9 blockage of an artery(s) and critical reduction of perfusion to the supply area that  
10 cannot be compensated by the collateral circulations (Bang et al., 2015). After  
11 blockage of the artery, the perfusion-compromised area undergoes necrosis  
12 (Kumar et al., 2010). Then, a highly interactive process of inflammation and repair  
13 system is activated, as evidenced by infiltration of inflammatory cells, along with  
14 induction of angiogenesis, neurogenesis and synaptogenesis (Ceulemans et al.,  
15 2010). Several reports have demonstrated that post-stroke neurogenesis depends  
16 significantly on neovascularization, suggesting that these processes work in  
17 concert to orchestrate a neurological recovery (Ohab et al., 2006; Thored et al.,  
18 2007; Xiong et al., 2011). Furthermore, histological data of stroke patients showed  
19 the importance of angiogenesis and vascular remodeling in the ischemic area, and  
20 higher blood vessel density in the stroke area indicates a better prognosis  
21 (Krupinski et al., 1994). Also, the therapy that increases angiogenesis has been  
22 shown to be beneficial in animal models of stroke (Sun et al., 2003). Hence,

1 angiogenesis in the stroke condition might be a good target for the therapy of the  
2 disease.

3         Angiogenesis is an important process for forming new blood vessels,  
4 manifested by extensive interactions between a wide variety of molecules including  
5 growth factors, cytokines, adhesion molecules, chemokines, enzymes and  
6 activators and inhibitors of angiogenesis (Carmeliet and Jain, 2011). In response to  
7 specific activators such as hypoxia or inflammation, angiogenesis regulators are  
8 expressed in the affected regions, causing the sprouting of vessels from existing  
9 one, and ultimately new vessels are formed (Carmeliet and Jain, 2011). Several  
10 angiogenesis regulators including VEGF,  $\beta$ FGF, angiopoietins and Tie2 are found  
11 to be expressed in stroke condition, suggesting that an angiogenic process is  
12 activated in the affected areas (Wang et al., 2002; Zhang and Chopp, 2015). This  
13 hypothesis is supported by the fact that new vessels are formed in the stroke area  
14 along with time dependent expression of angiogenesis regulators (Buga et al.,  
15 2014; Krupinski et al., 1994). Such process appears to be important for  
16 neurogenesis and neuroprotection as it was shown that new neurons are  
17 frequently found near the area of angiogenesis (Ohab et al., 2006; Thored et al.,  
18 2007).

19         Recently stem cell transplantation-based therapies are gaining much  
20 interest because of their ability to replace damaged neurons, neuroprotection,  
21 immunomodulation and angiogenesis (Chen et al., 2003; Chen et al., 2004; Sheikh  
22 et al., 2011). Among the stem cell types used for stroke therapy, mesenchymal

1 stem cells might be important because of its ability to produce growth factors,  
2 induce neuronal differentiation and modulate neuroinflammation (Nagai et al.,  
3 2007; Sheikh et al., 2011; Wakabayashi et al., 2010). In our previous study, we  
4 have found that a mesenchymal stem cell line transplantation increased VEGF  
5 mRNA expression along with many other growth factors in vivo in a MCAO rat  
6 model (Wakabayashi et al., 2010). Moreover, the analysis of MSC secretome  
7 shows that it can produce several angiogenic factors (Bakopoulou et al., 2015).  
8 Therefore, we hypothesized that MSC could modulate the angiogenesis system in  
9 stroke condition. To test this hypothesis, we transplanted a mesenchymal stem cell  
10 line in a rat stroke model and examined angiogenesis and the regulation of its  
11 underlying mechanism. We found that the transplantation increased vessel density  
12 in the stroke area possibly through increased expression of HIF-1 $\alpha$  and VEGF.

13

#### 14 **Materials and methods:**

##### 15 **Cell culture:**

16 Bone marrow cells were isolated from human fetal spinal vertebrae, and  
17 immortalized by introducing *v-Myc* oncogene using a retroviral vector (Nagai et al.,  
18 2007). A clone (B10) of immortalized bone marrow cells, which showed similar  
19 morphological and expressional phenotype and differentiation potentials as primary  
20 human mesenchymal stem cells, was expanded. This human mesenchymal stem  
21 cell line was cultured in complete MF<sup>®</sup> medium (Toyobo, Osaka, Japan) containing  
22 1% FCS and growth factor supplement (Wakabayashi et al., 2010).

1 Human primary microglia were isolated from human fetal brain and  
2 immortalized by transfecting a retroviral vector encoding v-*Myc*, as described  
3 previously (Nagai et al., 2001). A clone (HMO6), which shows similar  
4 morphological and expressional phenotype as primary microglia, was expanded.  
5 HMO6 was cultured in Dulbecco's modified Eagle medium (DMEM, Gibco,  
6 Invitrogen, Carlsbad, CA) of high glucose concentration, supplemented with 5%  
7 fetal bovine serum (Gibco), L-glutamine and antibiotics (Gibco) (Nagai et al., 2001).  
8 During cytokine treatment, 0.5% FBS containing DMEM was used for both B10 and  
9 HMO6 cells. For oxygen glucose deprivation (OGD) experiments, cells were  
10 incubated in an incubator in humid condition at 37°C, with 0.1% O<sub>2</sub> and 5% CO<sub>2</sub> for  
11 4 h. The composition of OGD medium was: NaCl 116 mM, KCl 5.4 mM, MgSO<sub>4</sub> 0.8  
12 mM, KH<sub>2</sub>PO<sub>4</sub> 0.44 mM, CaCl<sub>2</sub> 1.2 mM, NaHCO<sub>3</sub> 20 mM, Na<sub>2</sub>HPO<sub>4</sub> 0.33 mM, and  
13 Phenol Red 10 mg/l.

14 To investigate about the effect of B10 on microglia, a co-culture system of  
15 HMO6 and B10 was developed. HMO6 cells were cultured in a well of 6-well cell  
16 culture plate and B10 cells were in a cell culture insert (Millipore, Billerica, MA).  
17 After confluency, B10 cell containing cell culture insert was placed in the well of  
18 HMO6, resulting free movement of cell culture secreting molecules through the  
19 pores. However, there was no physical contact between B10 and HMO6 cells.  
20 DMEM medium containing 0.5% FBS was added and co-culture was continued up  
21 to 36 h.

22 Silencing IL-1 $\beta$  in B10 cells:

1 To silence human IL-1 $\beta$  mRNA expression, gene-specific silencing siRNA (Qingen,  
2 Valencia, CA) was transfected into B10 cells using HiPerfect transfection reagent  
3 (Qiagen) according to the manufacturer's protocol after optimizing the condition.  
4 For transfection, 10 nM siRNA and 3  $\mu$ l HiPerfect transfection reagent was used for  
5 1 well of 24-well plate. When transfection was done in 100 mm dishes, the amount  
6 of Hiperfect reagent was increased proportionate to surface area. Forty-eight hours  
7 after transfection, cell culture medium was changed with 0.5% FBS containing  
8 DMEM and cultured for further 24 h. Then the silencing effect was determined by  
9 real-time PCR and ELISA.

#### 10 **Animal model of focal ischemia**

11 Animal care was done according to the guideline of the experimental  
12 institute of Shimane University. All animals were kept under a constant  
13 temperature ( $23 \pm 2^\circ\text{C}$ ) and a light-dark cycle of 12 h. The animals were fed with  
14 commercially available normal rat diet and tap water *ad libitum*. All experimental  
15 protocol and procedures were approved by the Ethical Committee of Shimane  
16 University School of Medicine, and were done following the guideline of the  
17 experimental animal institute of Shimane University. Adult male Wister rats  
18 (Charles River, Yokohama, Japan), weighing 250 to 300 g, were used in this study.  
19 Transient focal cerebral ischemia was induced following an established method  
20 described previously (Wakabayashi et al., 2010). Briefly, rats were anesthetized  
21 with 4% halothane. Then the common, external and internal carotid arteries were  
22 exposed through a ventral midline incision. A silicon coated 4-0 monofilament

1 nylon suture with rounded tip was inserted through the right common carotid artery,  
2 and advanced until it occluded the middle cerebral artery. Then the anesthesia was  
3 reversed, and the rats were kept in a cage. After 90 min of occlusion, the rat was  
4 re-anesthetized and the nylon suture was withdrawn. Rectal temperature was  
5 maintained around 37°C throughout the surgical procedure using a feedback-  
6 regulated heating system.

### 7 **Intravenous injection of B10 human mesenchymal stem cells**

8 One day after MCAO, the rats were neurologically evaluated using a  
9 previously described neurological severity scoring system (Wakabayashi et al.,  
10 2010). Rats of similar neurological severity score were randomly divided into 3  
11 groups (n=5 for each time point and experiment type) to receive B10-cell  
12 transplantation, IL-1 $\beta$ -silenced B10-cell transplantation or phosphate buffered  
13 saline (PBS). IL-1 $\beta$  silenced B10 cells were transplanted 24 h after transfection of  
14 siRNA. For transplantation, a rat was anesthetized with 4% halothane, the jugular  
15 vein was exposed, and 3  $\times$  10<sup>6</sup> cells in 100  $\mu$ l of PBS or PBS-alone was injected  
16 intravenously (Wakabayashi et al., 2010). The tissue of a total 60 MCAO rats (25  
17 for PBS-treated groups, 25 for B10 transplanted group and 10 for IL-1 $\beta$  silenced  
18 B10 transplanted group) was used in this study.

### 19 **Immunohistochemical analysis**

20 Three, seven and 14 days after MCAO, rats were deeply anesthetized with  
21 isoflurane, and perfused transcardially with normal saline followed by 4%  
22 paraformaldehyde in 0.1 M phosphate buffer to fix the tissue. Then the brains were

1 removed, postfixed with same fixative, cryoprotected with 30% sucrose, and tissue  
2 blocks of 2 mm thickness were sectioned. For staining, a tissue slice of 10  $\mu$ m  
3 thickness was sectioned on a cryostat. After quenching endogenous peroxidase  
4 activity, the sections were incubated in a blocking solution containing 10% normal  
5 goat or horse serum and 0.2% Triton X-100 in PBS. Then the sections were  
6 incubated with anti-von-willebrand factor (vWF) IgG (rabbit, 1:100, Dako,  
7 Carpinteria, CA), anti-endoglin (CD105) IgG (goat, 1:200, R&D, Minneapolis, MN)  
8 anti-Iba-1 IgG (rabbit, 1:200, Wako, Richmond, VA), anti-HIF-1 $\alpha$  IgG (Mouse, 1:200  
9 Santa Cruz, Santa Cruz, CA), anti GFAP IgG (rabbit, 1:200, Dako), anti-p300 IgG  
10 (rabbit, 1:200, Santa Cruz), anti-VEGF IgG (goat, 1:100, Santa Cruz), anti-TGF $\beta$   
11 IgG (rabbit, 1:100, Dako), anti-IL-1 $\beta$  (goat, 1:200, Santa Cruz) and anti-human  
12 nuclei IgG (mouse, 1:100, Millipore). In the case of immunofluorescence staining,  
13 the tissue sections were then incubated with Texas red-conjugated or FITC-  
14 conjugated species-specific IgG, and nuclei were stained with Hoechst. For light  
15 microscopy, the section was incubated with biotin-conjugated species-specific IgG  
16 (1:100, Vector, Ingold Road, CA), followed by incubation with an avidin–biotin–  
17 peroxidase complex (ABC, Vector). The immune reaction products were visualized  
18 with 3, 30-diaminobenzidine (DAB, Sigma, St. Louis, MO), and the tissue was  
19 counterstained with Haematoxylin. Stained sections were examined under a  
20 fluorescent microscope (NIKON, ECLIPSE E600). For counting the cells, 3 tissue  
21 sections of 2 mm apart were stained, and cells were randomly counted in 5  
22 microscopic field of designated area at X400 magnification in a blinded manner.

1 The average cell number of a total 15 fields represented the cell number in that  
2 area of the rat brain. To determine vessel density in 3 and 7 days MCAO models, 3  
3 tissue sections of 2 mm apart were immunostained with vWF, photomicrographs of  
4 5 random microscopic fields at X400 magnification in the core and IBZ regions  
5 were taken, and analyzed the immunostained area by ImageJ software. The vessel  
6 density was calculated as percent of immunostained area occupied in a  
7 microscopic field. The average percent area of a total 15 fields represented the  
8 vessel density in that area of the rat brain

#### 9 **Total RNA isolation, reverse transcription and quantitative real time PCR**

10 Total RNA was isolated from cultured cells after appropriate treatment, or  
11 from infarct core, cortical ischemic border zone (IBZ) or contralateral cortex of  
12 PBS-treated or B10-transplanted rat brains 3 days after MCAO, using Trizol  
13 reagent (Invitrogen) according to the manufacturer's instructions. To prepare first  
14 strand cDNA, 2 µg of total RNA was reverse transcribed with reverse transcriptase  
15 enzyme (RiverTraAce, Toyobo, Osaka, Japan) in a 20 µl reaction mixture. To  
16 analyze mRNA level, real time PCR was performed with SyBr green PCR system  
17 (Power SyBr green, Applied Biosystem, Warrington, UK) using an ABI Prism 7300  
18 Sequence Detector system (Applied Biosystems). To detect mRNA of rat origin,  
19 real time PCR primers were designed using sequences of rat mRNA regions that  
20 are not in homology with human mRNA sequences of same gene (Wakabayashi et  
21 al., 2010). The mRNA level was normalized by corresponding GAPDH mRNA and  
22 quantified using relative quantification method.

## 1 **Western blot analysis**

2 Total protein was isolated from cultured cells after appropriate treatment, or  
3 from infarct core, IBZ or contralateral cortex of PBS-treated or B10-transplanted rat  
4 brain 3 days after MCAO using ice cold RIPA buffer (PBS, pH 7.4, 1% Nonidet p-  
5 40, 0.5% sodium deoxycholate, 0.1%SDS, 10 mg/ml PMSF, and 1 mg/ml aprotinin).  
6 To homogenize brain tissue, 20x wt/vol of RIPA buffer was used. Twenty to 60 µg  
7 of total protein was separated by SDS polyacrylamide gel electrophoresis,  
8 transferred to a PVDF membrane (Millipore, Billerica, MA). After blocking, the  
9 membrane was incubated with anti-HIF-1α (mouse, Santa Cruz), anti-p300 (rabbit,  
10 Santa Cruz), anti-VEGF (goat, Santa Cruz) or anti-TGFβ (rabbit, Dako) IgG. Then  
11 the membrane was incubated with infrared (IR) fluorophore-conjugated species-  
12 specific IgG (Li-Cor, Lincoln, NE). Immunoreactive proteins in the membrane were  
13 detected using an IR scanner (Li-Cor) according to the manufacturer's protocol.

## 14 **ELISA.**

15 Culture supernatants were collected from the cell cultures, and the  
16 concentration of IL-1β was measured by sandwich ELISA using a human IL-1β  
17 platinum ELISA Kit (Invitrogen) according to the manufacturer's instructions. Briefly,  
18 culture supernatants were centrifuged at 14,000 rpm for 10 min at 4°C to remove  
19 cell debris. Equal volume of sample diluent, provided by the manufacturer, was  
20 added to the culture supernatant. Then 100 µl of diluted samples were added to  
21 the wells of a 96-well plate coated with anti-IL-1β antibody and incubated at room  
22 temperature for 2 h. After wash, a biotin conjugated anti-human IL-1β antibody

1 (100  $\mu$ l) was added to the wells and incubated for 2 h at room temperature. Then  
2 HRP-conjugated streptavidin (100  $\mu$ l) was added to the wells and incubated for 1 h  
3 at room temperature. After wash, TMB substrate (100  $\mu$ l) was added to the wells  
4 and incubates at room temperature in a dark condition for 10 min. The reaction  
5 was terminated by adding a stop solution provided by the manufacturer, and optical  
6 densities were measured with a plate reader (EAR400, SLT-Lab instruments). IL-  
7 1 $\beta$  concentration was calculated after preparing a standard curve using human IL-  
8 1 $\beta$  standard (Invitrogen).

#### 9 **Statistical analysis**

10 The numerical data are presented as mean  $\pm$  SD. Statistical differences among the  
11 groups were assessed by one-way ANOVA followed by Scheffe's post hoc test, or  
12 paired t-test. The statistical significance level was set at  $p < 0.05$

13

#### 14 **Result:**

15 **B10 transplantation increased angiogenesis in a rat MCAO model.** In a  
16 previous report, we have shown that B10 transplantation increased VEGF mRNA  
17 expression in a rat MCAO model after 3 days (Wakabayashi et al., 2010). Hence,  
18 we chose to investigate angiogenesis starting at that time point. Staining of blood  
19 vessels with vWF antibody revealed that in PBS-treated MCAO rats, the vessel  
20 density was significantly decreased in the core region compared to ischemic border  
21 zone (IBZ) at 3 days (Figure 1A and 1B). Compared to corresponding areas of  
22 PBS-treated group, B10-transplanted group showed increased vessel density in

1 both ischemic core and IBZ region in the cortical areas at 3 days (Figure 1A and  
2 1B). At 7 days after MCAO, the difference still persisted in the core region; however,  
3 the vessel density was similar in the IBZ area of PBS-treated and B10-transplanted  
4 groups (Figure 1C and 1D). Similar results were observed at 14 days after MCAO,  
5 where the difference of vessel density was found mainly in the core region  
6 (Supplemental Figure 1).

7 **Transplanted B10 cells localized in the ischemic area and increased newly**  
8 **formed vessels in rat MCAO model.** Next, we analyzed the distribution of  
9 transplanted B10 cells in MCAO rat brains. Since it is a human origin mesenchymal  
10 stem cell line, we used antibody against human nuclear antigen (HuN) to identify  
11 transplanted B10 cells in rat host brains. As we have seen in our previous study  
12 (Wakabayashi et al., 2010), B10 cells mainly accumulated in the core and IBZ  
13 regions of the brains at 3 days after MCAO (Figure 2A). Seven days after MCAO,  
14 no B10 cells were detectable in the rat brains (Figure 2B).

15 To analyze further about the effects of B10 transplantation on angiogenesis  
16 in MCAO condition, we evaluated newly formed vessels by endoglin  
17 immunostaining. Endoglin is suggested to be a better marker for newly formed  
18 vessels, whereas vWF can detect both newly formed and normal vessels (Yao Y et  
19 al., 2005). Immunostaining results showed that at 3 days after MCAO, endoglin  
20 positive cell number was increased in both core and IBZ areas of B10 transplanted  
21 rats (Figure 2C and 2E). At 7 days after MCAO, endoglin stains adopted  
22 microvessel-like structures in the core and IBZ regions of both PBS and B10

1 transplanted rats (Figure 2D). Moreover, some endoglin positive cells were present  
2 along with microvessel-like structure in B10 transplanted rats, especially in the core  
3 area. The average diameter of those microvessel-like structures were 23.6  $\mu\text{m}$   
4 (range 15 – 31.2  $\mu\text{m}$ ) and 25.9  $\mu\text{m}$  (range 14.4 – 40.5  $\mu\text{m}$ ) in the core region of  
5 PBS-treated and B10 transplanted MCAO rats, respectively. In the IBZ region of  
6 PBS-treated and B10 transplanted rats, the average diameter was 31  $\mu\text{m}$  (range  
7 20 – 41.7  $\mu\text{m}$ ) and 28.3  $\mu\text{m}$  (range 18.7 – 40.8  $\mu\text{m}$ ), respectively. Counting of these  
8 microvessel-like structures revealed that the number was significantly increased in  
9 the core region of B10 transplanted rats (Figure 2F). At 14 days after MCAO,  
10 endoglin area was still increased in the core region of B10-transplanted rats. In the  
11 IBZ region of both PBS-treated and B10-transplanted rats, well-formed  
12 microvessels were seen (Supplemental figure 2).

13 **B10 transplantation affected the mRNA expression of angiogenesis-related**  
14 **genes in rat MCAO model.** Next, we have checked mRNA expression of several  
15 angiogenesis-related factors in MCAO rat brains. The results revealed that the  
16 mRNA levels of angiopoetin2 (Ang2), placenta growth factor (PIGF) and Tie1 were  
17 considerably increased in PBS-treated MCAO rats brains, whereas the mRNA of  
18 angiopoetin1 (Ang1), HIF-1 $\alpha$  and EGL-Nine homologs (EGLNs) were not increased  
19 much (Figure 3). Compared to PBS-treated group, the mRNA levels of Ang1, HIF-  
20 1 $\alpha$  and PIGF were significantly increased in the IBZ region of B10-transplanted  
21 group. In the core region of B10-transplanted group, the mRNA of HIF-1 $\alpha$  and  
22 PIGF were increased. Conversely, compared to PBS-treated group, the expression

1 of Ang2, Flt1, Tie1 and Tie2 mRNA was decreased in IBZ region of B10-  
2 transplanted group. However, the mRNA levels of EGLNs, the enzymes that  
3 regulate hypoxia-dependent stability of HIF-1 $\alpha$ , were similar between 2 groups in  
4 all areas that we have checked.

5 **B10 transplantation increased HIF-1 $\alpha$  protein level in a rat MCAO model.** Next,  
6 we investigated about protein expression of HIF-1 $\alpha$ . Western blotting results  
7 demonstrated that HIF-1 $\alpha$  protein was barely detectable in the core, IBZ or  
8 contralateral cortical areas of PBS-treated MCAO rat brains 3 days after MCAO  
9 (Figure 4A). However, a significant increase of HIF-1 $\alpha$  protein was seen mainly in  
10 the core and IBZ region of B10-treated group (Figure 4A). On the other hand, the  
11 level of p300, a coactivator of HIF-1 $\alpha$ , was not changed (Figure 4A). To analyze  
12 their localizations in B10-transplanted rat brain tissue, double immunofluorescence  
13 staining of HIF-1 $\alpha$  and GFAP (astrocyte marker), or HIF-1 $\alpha$  and Iba-1  
14 (macrophage/microglia marker) were done. We did not find GFAP positive  
15 astrocytes in the core region of rat brains 3 days after MCAO, where Iba-1 positive  
16 macrophage/microglia were detected. In the IBZ region, both GFAP positive  
17 astrocytes and Iba-1 positive macrophage/microglia were detectable. In the core  
18 region, HIF-1 $\alpha$  was expressed mainly in Iba-1 positive macrophage/microglia  
19 (Figure 4B). In the IBZ region, HIF-1 $\alpha$  was found to be expressed in both GFAP  
20 positive astrocytes and Iba-1 positive macrophage/microglia (Figure 4B and 4C).  
21 As p300 plays a role in HIF-1 $\alpha$  activation (Ruas et al., 2010), we checked its co-

1 localization with HIF-1 $\alpha$ . The immunofluorescence results showed that indeed,  
2 p300 was co-localized with HIF-1 $\alpha$  (Figure 4D).

### 3 **B10 transplantation increased VEGF protein level in a rat MCAO model.**

4 VEGF is a potent angiogenesis factor. As VEGF is a downstream target of HIF-1 $\alpha$   
5 (Forsythe et al., 1996), we checked its regulation by B10 transplantation in MCAO  
6 rat model. Western blotting results demonstrated that VEGF protein was detectable  
7 in the core, IBZ and contralateral cortices of PBS-treated rat brains 3 days after  
8 MCAO, which was increased in those areas in B10-transplanted rats (Figure 5A).  
9 Immunostaining results showed that in the core and IBZ regions, VEGF expressed  
10 mainly in round shaped cells (Figure 5B). Double immunofluorescence staining  
11 showed that VEGF expressed mainly in Iba-1 positive macrophage/microglia in  
12 both core and IBZ region of PBS-treated and B10-transplanted rats (Figure 5C and  
13 5D).

### 14 **B10 transplantation increased TGF $\beta$ production in a rat MCAO model.**

15 Previous reports have shown that TGF $\beta$  plays an important role in the development  
16 of blood vessels (Nguyen et al., 2011). Hence, we checked regulation of TGF  
17 production in MCAO rat model. Western blotting results showed that TGF $\beta$  protein  
18 level is undetectable in the contralateral cortex of the rats, whereas the protein was  
19 barely detectable in the core and IBZ cortical areas in PBS-treated rats 3 days after  
20 MCAO (Figure 6A). The level of TGF $\beta$  was increased in the core and IBZ region in  
21 B10-transplanted rats (Figure 6A). Immunofluorescence staining also confirmed

1 the result, showing an increase of TGF $\beta$  expression in the core and IBZ region of  
2 B10-transplanted rats (Figure 6B).

3 **Regulation of VEGF expression in macrophage/microglia by B10.** We  
4 investigated about the regulation of VEGF expression in macrophage/microglia by  
5 B10 cells using an in vitro cell culture system. In previous reports, it has been  
6 shown that inflammatory cytokines affect angiogenesis through production of  
7 VEGF (Asano-Kato et al., 2005). Our real-time PCR result showed that stimulation  
8 of a macrophage/microglia cell line (HMO6) or B10 with IL-1 $\beta$  time-dependently  
9 increased VEGF mRNA level (Figure 7A and 7B). However, the peak of VEGF  
10 mRNA expression was 8 h in B10 and 24 h in HMO6 (Figure 7A and 7B). Next, we  
11 checked the effects of B10 on HMO6 in respect of IL-1 $\beta$  and VEGF expression.  
12 Here, we used a co-culture system using cell culture inserts, so that cell products  
13 can affect the other cell type without any contact signaling. The real-time PCR  
14 result showed that IL-1 $\beta$  mRNA level was very high in B10 cells in respect to  
15 HMO6 (Figure 6C) although at protein level, the expression was similar among the  
16 cell types (Supplemental Figure 3). In co-culture system, IL-1 $\beta$  mRNA was  
17 increased in B10 cells, whereas it was decreased in HMO6 after 36 h (Figure 7C).  
18 However, VEGF mRNA level was increased in co-culture system in both cell types  
19 (Figure 7D).

20 Next, we investigated about the effect of B10 cells on HIF-1 $\alpha$  and VEGF  
21 proteins expression in HMO6 in oxygen-glucose deprivation (OGD) and non-OGD  
22 conditions. Total cell lysate of HMO6 was isolated after co-culturing with B10 cells

1 for 24 h. Western blotting results showed that compared to HMO6 native culture,  
2 VEGF protein level was increased in HMO6 co-cultured with B10 cells whereas  
3 HIF-1 $\alpha$  protein level was not affected (Figure 6E and 6F, non-OGD). In OGD  
4 condition, both HIF-1 $\alpha$  and VEGF were increased similarly in HMO6 native culture  
5 and co-cultured with B10 cells (Figure 7E and 7F).

6 **B10 transplantation regulated IL-1 $\beta$  production in a rat MCAO model.** Next,  
7 we checked IL-1 $\beta$  expression *in vivo* in rat MCAO models. Our immunostaining  
8 results showed that IL-1 $\beta$  producing cells were found mainly in the core region of  
9 rats 3 days after MCAO, with some cells having low level of IL-1 $\beta$  at the IBZ  
10 (Figure 8A and 8B). Counting of the immunoreactive cells revealed that the number  
11 of IL-1 $\beta$  positive cells was increased in B10-transplanted group (Figure 8C). To  
12 identify the cells that produced IL-1 $\beta$ , we employed double immunofluorescence  
13 staining. The staining results showed that in PBS-treated rat MCAO model, IL-1 $\beta$  is  
14 expressed mainly in ED-1 positive macrophage/microglia (Figure 8D). In B10-  
15 transplanted group, IL-1 $\beta$  was expressed in ED-1 positive macrophage/microglia as  
16 well as in human nuclear antigen positive transplanted B10 cells (Figure 8D and  
17 8E).

18 **Silencing IL-1 $\beta$  in B10 decreased its ability to induce endoglin in a rat MCAO**  
19 **model.** To explore further about the role of IL-1 $\beta$  in B10-induced angiogenesis, we  
20 silenced its expression in B10 cells by mRNA specific siRNA transfection. IL-1 $\beta$   
21 mRNA specific siRNA decreased the expression both at mRNA and protein level to  
22 less than 40% (Supplemental Figure 4). These IL-1 $\beta$ -silenced B10 cells were

1 transplanted in rat MCAO model. Immunostaining results showed that endoglin  
2 positive area was significantly decreased in the core and IBZ regions of IL-1 $\beta$ -  
3 silenced B10 transplanted rats compared to non-silenced B10 transplanted rats  
4 both at 3 days (Figure 9A and 9B) and 7 days (Figure 9C and 9D) after MCAO.  
5 However, endoglin positive vessel-like structure appeared early in IL- $\beta$ -silenced  
6 B10 transplanted rat brains, as early as 3 days after MCAO, which was not  
7 observed in non-silenced B10 transplanted condition (Figure 9A and 9C).

#### 8 **Discussion:**

9       Vascular remodeling and angiogenesis plays a key role in the pathological  
10 outcome of stroke condition (Krupinski et al., 1994; Lapi and Colantuoni, 2015;  
11 Zhang and Chopp, 2015). Experimental evidences suggested that a better vascular  
12 remodeling system improved not only the blood supply, but also neuronal function  
13 (Lapi and Colantuoni, 2015). This fact is supported by the clinical observations that  
14 the stroke patients having better local circulation attain better outcome (Krupinski  
15 et al., 1994). In our previous study, we have demonstrated that B10 transplantation  
16 improves the neurological performances of a rat MCAO model (Wakabayashi et al.,  
17 2010). In that study, we have found that the mRNA expression of VEGF-A was  
18 increased in MCAO rat brains at an earlier time point than other growth factors and  
19 cytokines. Importantly, compared to control, B10 transplantation increased VEGF-  
20 A mRNA in MCAO rat brains at that time point. Consistent with that finding, in this  
21 study we have demonstrated that B10 transplantation increased vessel density in  
22 the core and IBZ region from 3 days after MCAO. Besides, endoglin staining was

1 increased in B10 transplantation condition, and the difference with PBS-treated  
2 MCAO rats was persisted after the disappearance of transplanted cells, signifying  
3 a lasting effect of transplantation on angiogenesis. Hence, early regulation of  
4 angiogenesis could be an important feature of B10 transplantation-mediated  
5 neurological improvement in MCAO condition.

6 VEGF family proteins including VEGF-A and PlGF are considered to be  
7 potent angiogenic factors (Holmes and Zachary, 2005). Here, we have  
8 demonstrated VEGF-A at protein level and PlGF mRNA were increased by B10  
9 transplantation, suggesting a central role of VEGF family proteins in MSC-  
10 mediated angiogenesis in cerebral ischemic condition. However, VEGF not only  
11 increases angiogenesis, but also reported to increase the vascular permeability  
12 leading to cerebral edema (Zhang et al., 2000). Our real-time PCR results showed  
13 that along with VEGF, B10 transplantation increased angiopoietin1 (Ang1)  
14 expression in IBZ region. Ang1 is considered to stabilize newly formed vessels  
15 through the activation of Tie2 receptor during angiogenesis (Brindle et al., 2006;  
16 Fiedler and Augustin, 2006). Ang1-mediated Tie2 activation is reported to be  
17 antagonized by Ang2 (Maisonpierre et al., 1997; Scharpfenecker, 2005).  
18 Interestingly the mRNA level of Ang2 was high in PBS group compared to B10  
19 transplanted MCAO rats in IBZ area. This finding indicates that Ang1-Tie2  
20 dependent new vessel stabilization could be well achieved in B10 transplanted  
21 condition. Moreover, TGF $\beta$  expression was increased in B10 transplanted MCAO  
22 rats. TGF $\beta$  is demonstrated to regulate angiogenesis in concert with VEGF, and it

1 is also shown that abrogation of the interaction of VEGF and TGF $\beta$  caused  
2 abnormal vessel formation (Ferrari et al., 2009). Hence, B10 transplantation might  
3 regulate the process of angiogenesis along with stabilization of newly formed  
4 vessels, and provides better perfusion efficiency.

5 In ischemic condition, VEGF expression is mainly regulated by HIF-1 $\alpha$ , a  
6 hypoxia regulated transcription factor (Chen et al., 2010). In our study, we found  
7 that HIF-1 $\alpha$  mRNA as well as protein was increased in B10 transplanted rat brains.  
8 In normoxia, HIF-1 $\alpha$  becomes a target of proteasome system by prolyl  
9 hydroxylases and rapidly degraded (Salceda and Caro, 1997). In hypoxic condition,  
10 the activity of prolyl hydroxylase is inhibited, causing stabilization of HIF-1 $\alpha$   
11 (Jaakkola et al., 2001). Although we did not investigate about the enzyme activity,  
12 the mRNA expression of prolyl hydroxylases EGLN1, EGLN2 and EGLN3 was not  
13 changed in B10 transplanted rats. In OGD condition, HIF-1 $\alpha$  protein level is known  
14 to increase due to decreased activity of prolyl hydroxylases (Souvenir et al., 2014).  
15 In our culture condition, HIF-1 $\alpha$  protein level in native cultured and B10 co-cultured  
16 HMO6 were similarly increased after OGD, indicating that B10 did not affect prolyl  
17 hydroxylase enzyme activity in HMO6. Therefore, B10 transplantation might not  
18 increase HIF-1 $\alpha$  protein through inhibiting EGLNs, rather by regulating its  
19 transcription.

20 In B10 transplanted rats, we have found that both HIF-1 $\alpha$  and VEGF were  
21 expressed in the core and IBZ regions. VEGF is reported to be express in  
22 astrocytes, neurons and microglia (Greenberg and Jin, 2013). However, soon after

1 stroke, microglia are considered to be the main source of VEGF (Greenberg and  
2 Jin, 2013). In this study, we demonstrated that although HIF-1 $\alpha$  expressed both in  
3 astrocytes and macrophage/microglia, VEGF was rarely found to be positive in  
4 astrocytes during this early time point, indicating the time-dependent role of  
5 different cell-types in angiogenesis (Greenberg and Jin, 2013). In a previous study,  
6 we have found that B10 transplantation decreased total number of  
7 macrophage/microglia in the core and IBZ regions of MCAO rats (Sheikh et al.;  
8 2011), suggesting a possibility that total VEGF expression might be decreased by  
9 B10 transplantation. However, microglia of M2 phenotype are demonstrated to  
10 express VEGF (Lamagna et al., 2006). In our previous studies (Sheikh et al., 2011;  
11 Wakabayashi et al., 2010), we showed that B10 transplantation not only increased  
12 VEGF, IL-10 and IL-4, but also decreased iNOS. These facts might imply that  
13 although it decreases the total macrophage/microglia cell number, it influences  
14 their M2 transition, and thereby increases VEGF and angiogenic factors expression.  
15 Such M2 transition of macrophage/microglia could be important for early  
16 angiogenesis and reparative processes, and subsequent neurological improvement  
17 that are seen in B10-transplanted condition.

18 As inflammation and angiogenesis are intimately related (Asano-Kato et al.,  
19 2005; Parmeggiani et al., 2010), we focused to investigate whether MSC  
20 transplantation modulates angiogenesis process through the regulation of  
21 inflammatory cytokines. Our real-time PCR and ELISA results showed that IL-1 $\beta$   
22 expression was high in B10 cells. Also, IL-1 $\beta$  time-dependently increased VEGF

1 mRNA expression in both B10 and a human microglia cell line (HMO6).  
2 Interestingly, co-culture of B10 and HMO6 increased IL-1 $\beta$  in B10, but decreased it  
3 in HMO6 cells. In a previous study, we showed that B10 transplantation increased  
4 IL-1 $\beta$  mRNA slightly in the core region at 3 days, and significantly in the IBZ region  
5 at 7 days after MCAO. In this study, IL-1 $\beta$  expressing cells were found to be  
6 modestly, but significantly increased in B10 transplanted rats 3 days after MCAO,  
7 where many of the IL-1 $\beta$  expressing cells were transplanted B10. These results are  
8 suggesting that B10 could be an important source of IL-1 $\beta$  in B10 transplanted  
9 MCAO rats, which increases endothelial cell proliferation and angiogenesis in the  
10 animal model (Yang et al., 2012; Voronov et al., 2003). The results of silencing  
11 study proved that hypothesis, since inhibiting expression of IL-1 $\beta$  in B10 cells  
12 decreases endoglin positive area in MCAO rat brains. Interestingly, vessel-like  
13 structure appeared early in IL-1 $\beta$ -silenced B10 transplanted rat brains, suggesting  
14 that although it increases endothelial cell proliferation, IL-1 $\beta$  might have a negative  
15 effect on vessel-like structure formation. In addition to its effects on proliferation,  
16 previous studies have demonstrated a role of IL-1 $\beta$  on endothelial cell apoptosis  
17 (Rivera et al., 2013; Zhu et al., 2014). Hence, a controlled expression of IL-1 $\beta$   
18 could be important, where an endothelial proliferation, apoptosis, arrangement to  
19 vessel-like structure and maturation are balanced for proper angiogenesis. It will be  
20 interesting to investigate further the role of IL-1 $\beta$ , and its interaction with the factors  
21 expressed by B10 cells that mediated angiogenesis in vivo in cerebral ischemic  
22 condition.

1           In conclusion, our findings provide a new insight of the mechanism of  
2 angiogenesis in cerebral ischemia condition, its regulation by mesenchymal stem  
3 cell transplantation and the importance of IL-1 $\beta$  on that regulation. Such  
4 understanding could be valuable for the improvement of stroke therapy.

5  
6 **Conflict of interest:** All authors declare no conflict of interest associated with this  
7 study.

8  
9 **References:**

- 10 Asano-Kato, N., Fukagawa, K., Okada, N., Kawakita, T., Takano, Y., Dogru, M.,  
11           Tsubota, K., Fujishima, H., 2005. TGF-beta1, IL-1beta, and Th2 cytokines  
12           stimulate vascular endothelial growth factor production from conjunctival  
13           fibroblasts. *Experimental Eye Research*. 80, 555-560.
- 14 Bélanger, M., Allaman, I., Magistretti, P.J., 2011. Brain energy metabolism: Focus  
15           on Astrocyte-neuron metabolic cooperation. In: *Cell Metabolism*. Vol. 14,  
16           ed.^eds., pp. 724-738.
- 17 Bakopoulou, A., Kritis, A., Andreadis, D., Papachristou, E., Leyhausen, G., Koidis,  
18           P., Geurtsen, W., Tsiftoglou, A., 2015. Angiogenic Potential and Secretome  
19           of Human Apical Papilla Mesenchymal Stem Cells in Various Stress  
20           Microenvironments. *Stem Cells and Development*. 24, 2496-2512.
- 21 Bang, O.Y., Goyal, M., Liebeskind, D.S., Najm, M., Rubiera, M., Fainardi, E., 2015.  
22           Collateral Circulation in Ischemic Stroke: Assessment Tools and  
23           Therapeutic Strategies. *Stroke*. 46, 3302-3309.
- 24 Brindle, N.P.J., Saharinen, P., Alitalo, K., 2006. Signaling and functions of  
25           angiopoietin-1 in vascular protection. In: *Circulation Research*. Vol. 98,  
26           ed.^eds., pp. 1014-1023.

- 1 Buga, A.M., Margaritescu, C., Scholz, C.J., Radu, E., Zelenak, C., Popa-Wagner,  
2 A., 2014. Transcriptomics of post-stroke angiogenesis in the aged brain.  
3 *Frontiers in Aging Neuroscience*. 6.
- 4 Carmeliet, P., Jain, R.K., 2011. Molecular mechanisms and clinical applications of  
5 angiogenesis. *Nature*. 473, 298-307.
- 6 Ceulemans, A.G., Zgavc, T., Kooijman, R., Hachimi-Idrissi, S., Sarre, S., Michotte,  
7 Y., 2010. The dual role of the neuroinflammatory response after ischemic  
8 stroke: modulatory effects of hypothermia. *J Neuroinflammation*. 7, 74.
- 9 Chen, C., Ostrowski, R.P., Zhou, C., Tang, J., Zhang, J.H., 2010. Suppression of  
10 hypoxia-inducible factor-1 $\alpha$  and its downstream genes reduces acute  
11 hyperglycemia-enhanced hemorrhagic transformation in a rat model of  
12 cerebral ischemia. *Journal of Neuroscience Research*. 88, 2046-2055.
- 13 Chen, J., Li, Y., Katakowski, M., Chen, X., Wang, L., Lu, D., Lu, M., Gautam, S.C.,  
14 Chopp, M., 2003. Intravenous bone marrow stromal cell therapy reduces  
15 apoptosis and promotes endogenous cell proliferation after stroke in female  
16 rat. *Journal of Neuroscience Research*. 73, 778-786.
- 17 Chen, J., Li, Y., Zhang, R., Katakowski, M., Gautam, S.C., Xu, Y., Lu, M., Zhang,  
18 Z., Chopp, M., 2004. Combination therapy of stroke in rats with a nitric oxide  
19 donor and human bone marrow stromal cells enhances angiogenesis and  
20 neurogenesis. *Brain Research*. 1005, 21-28.
- 21 Ferrari, G., Cook, B.D., Terushkin, V., Pintucci, G., Mignatti, P., 2009.  
22 Transforming growth factor-beta 1 (TGF-beta1) induces angiogenesis  
23 through vascular endothelial growth factor (VEGF)-mediated apoptosis.  
24 *Journal of cellular physiology*. 219, 449-458.
- 25 Fiedler, U., Augustin, H.G., 2006. Angiopoietins: a link between angiogenesis and  
26 inflammation. In: *Trends in Immunology*. Vol. 27, ed. eds., pp. 552-558.
- 27 Forsythe, J.A., Jiang, B.H., Iyer, N.V., Agani, F., Leung, S.W., Koos, R.D.,  
28 Semenza, G.L., 1996. Activation of vascular endothelial growth factor gene

1 transcription by hypoxia-inducible factor 1. *Molecular and cellular biology*. 16,  
2 4604-4613.

3 Greenberg, D.A., Jin, K., 2013. Vascular endothelial growth factors (VEGFs) and  
4 stroke. *Cellular and molecular life sciences : CMLS*. 70, 1753-1761.

5 Holmes, D.I.R., Zachary, I., 2005. The vascular endothelial growth factor (VEGF)  
6 family: angiogenic factors in health and disease. *Genome biology*. 6, 209.

7 Jaakkola, P., Mole, D.R., Tian, Y.M., Wilson, M.I., Gielbert, J., Gaskell, S.J., von  
8 Kriegsheim, A., Hebestreit, H.F., Mukherji, M., Schofield, C.J., Maxwell, P.H.,  
9 Pugh, C.W., Ratcliffe, P.J., 2001. Targeting of HIF- $\alpha$  to the von Hippel-  
10 Lindau ubiquitylation complex by O<sub>2</sub>-regulated prolyl hydroxylation. *Science*.  
11 292, 468-472.

12 Krupinski, J., Kaluza, J., Kumar, P., Kumar, S., Wang, J., 1994. Role of  
13 angiogenesis in patients with cerebral ischemic stroke. *Stroke*. 25, 1794-  
14 1798.

15 Kumar, V., Abbas, A.K., Fausto, N., Aster, J.C., 2010. Robbins and Cotran  
16 Pathologic Basis of Disease. Robbins and Cotran Pathologic Basis of  
17 Disease. 23-24.

18 Lamagna, C., Aurrand-Lions, M., Imhof, B.a., 2006. Dual role of macrophages in  
19 tumor growth and angiogenesis. *Journal of leukocyte biology*. 80, 705-713.

20 Lapi, D., Colantuoni, A., 2015. Remodeling of Cerebral Microcirculation after  
21 Ischemia-Reperfusion. In: *Journal of Vascular Research*. Vol. 52, ed.^eds.,  
22 pp. 22-31.

23 Liebeskind, D.S., 2003. Collateral circulation. In: *Stroke*. Vol. 34, ed.^eds., pp.  
24 2279-2284.

25 Maisonpierre, P.C., Suri, C., Jones, P.F., Bartunkova, S., Wiegand, S.J.,  
26 Radziejewski, C., Compton, D., McClain, J., Aldrich, T.H., Papadopoulos, N.,  
27 Daly, T.J., Davis, S., Sato, T.N., Yancopoulos, G.D., 1997. Angiopoietin-2, a  
28 Natural Antagonist for Tie2 That Disrupts in vivo Angiogenesis. *Science*.  
29 277, 55-60.

1 Nagai, a., Nakagawa, E., Hatori, K., Choi, H.B., McLarnon, J.G., Lee, M.a., Kim,  
2 S.U., 2001. Generation and characterization of immortalized human  
3 microglial cell lines: expression of cytokines and chemokines. *Neurobiology*  
4 *of disease*. 8, 1057-1068.

5 Nagai, A., Kim, W.K., Lee, H.J., Jeong, H.S., Kim, K.S., Hong, S.H., Park, I.H., Kim,  
6 S.U., 2007. Multilineage potential of stable human mesenchymal stem cell  
7 line derived from fetal marrow. *PLoS ONE*. 2.

8 Nguyen, H.-L., Lee, Y.J., Shin, J., Lee, E., Park, S.O., McCarty, J.H., Oh, S.P.,  
9 2011. TGF- $\beta$  signaling in endothelial cells, but not neuroepithelial cells, is  
10 essential for cerebral vascular development. *Laboratory investigation; a*  
11 *journal of technical methods and pathology*. 91, 1554-1563.

12 Ohab, J.J., Fleming, S., Blesch, A., Carmichael, S.T., 2006. A neurovascular niche  
13 for neurogenesis after stroke. *J Neurosci*. 26, 13007-13016.

14 Parmeggiani, F., Campa, C., Costagliola, C., Incorvaia, C., Sheridan, C., Semeraro,  
15 F., De Nadai, K., Sebastiani, A., 2010. Inflammatory mediators and  
16 angiogenic factors in choroidal neovascularization: Pathogenetic  
17 interactions and therapeutic implications. In: *Mediators of Inflammation*. Vol.  
18 2010, ed. eds.

19 Rivera, J.C., Sitaras, N., Noueihed, B., Hamel, D., Madaan, A., Zhou, T., Honoré,  
20 J.C., Quiniou, C., Joyal, J.S., Hardy, P., Sennlaub, F., Lubell, W., Chemtob,  
21 S., 2013. Microglia and interleukin-1 $\beta$  in ischemic retinopathy elicit  
22 microvascular degeneration through neuronal semaphorin-3A. *Arterioscler*  
23 *Thromb Vasc Biol*. 33, 1881-1891.

24 Ruas, J.L., Berchner-Pfannschmidt, U., Malik, S., Gradin, K., Fandrey, J., Roeder,  
25 R.G., Pereira, T., Poellinger, L., 2010. Complex regulation of the  
26 transactivation function of hypoxia-inducible factor-1 $\alpha$  by direct interaction  
27 with two distinct domains of the creb-binding protein/p300. *Journal of*  
28 *Biological Chemistry*. 285, 2601-2609.

1 Salceda, S., Caro, J., 1997. Hypoxia-inducible factor 1 $\alpha$  (HIF-1 $\alpha$ ) protein is rapidly  
2 degraded by the ubiquitin-proteasome system under normoxic conditions.  
3 Its stabilization by hypoxia depends on redox-induced changes. *Journal of*  
4 *Biological Chemistry.* 272, 22642-22647.

5 Scharpfenecker, M., 2005. The Tie-2 ligand Angiopoietin-2 destabilizes quiescent  
6 endothelium through an internal autocrine loop mechanism. *Journal of Cell*  
7 *Science.* 118, 771-780.

8 Sheikh, A.M., Nagai, A., Wakabayashi, K., Narantuya, D., Kobayashi, S.,  
9 Yamaguchi, S., Kim, S.U., 2011. Mesenchymal stem cell transplantation  
10 modulates neuroinflammation in focal cerebral ischemia: Contribution of  
11 fractalkine and IL-5. *Neurobiology of Disease.* 41, 717-724.

12 Souvenir, R., Flores, J.J., Ostrowski, R.P., Manaenko, A., Duris, K., Tang, J., 2014.  
13 Erythropoietin Inhibits HIF-1?? Expression via Upregulation of PHD-2  
14 Transcription and Translation in an In Vitro Model of Hypoxia-Ischemia.  
15 *Translational Stroke Research.* 5, 118-127.

16 Sun, Y., Jin, K., Xie, L., Childs, J., Mao, X.O., Logvinova, A., Greenberg, D.A.,  
17 2003. VEGF-induced neuroprotection, neurogenesis, and angiogenesis after  
18 focal cerebral ischemia. *Journal of Clinical Investigation.* 111, 1843-1851.

19 Thored, P., Wood, J., Arvidsson, A., Cammenga, J., Kokaia, Z., Lindvall, O., 2007.  
20 Long-term neuroblast migration along blood vessels in an area with  
21 transient angiogenesis and increased vascularization after stroke. *Stroke.*  
22 38, 3032-3039.

23 Voronov, E., Shouval, D.S., Krelin, Y., Cagnano, E., Benharroch, D., Iwakura, Y.,  
24 Dinarello, C.A., Apte, R.N., 2003. IL-1 is required for tumor invasiveness  
25 and angiogenesis. *Proc Natl Acad Sci U S A.* 100, 2645-2650.

26 Wakabayashi, K., Nagai, A., Sheikh, A.M., Shiota, Y., Narantuya, D., Watanabe, T.,  
27 Masuda, J., Kobayashi, S., Kim, S.U., Yamaguchi, S., 2010. Transplantation  
28 of human mesenchymal stem cells promotes functional improvement and

1 increased expression of neurotrophic factors in a rat focal cerebral ischemia  
2 model. *Journal of Neuroscience Research*. 88, 1017-1025.

3 Wang, M.M., Klaus, J.a., Joh, H.-D., Traystman, R.J., Hurn, P.D., 2002.  
4 Postischemic angiogenic factor expression in stroke-prone rats.  
5 *Experimental neurology*. 173, 283-288.

6 Xiong, Y., Mahmood, A., Chopp, M., 2011. Angiogenesis, neurogenesis and brain  
7 recovery of function following injury. *Curr Opin Investig Drugs*. 11, 298-308.

8 Yang, L., Guo, X.G., Du, C.Q., Yang, J.X., Jiang, D.M., Li, B., Zhou, W.J., Zhang,  
9 F.R., 2012. Interleukin-1 beta increases activity of human endothelial  
10 progenitor cells: involvement of PI3K-Akt signaling pathway. *Inflammation*.  
11 35, 1242-1250.

12 Yao, Y., Kubota, T., Takeuchi, H., Sato, K., 2005. Prognostic significance of  
13 microvessel density determined by an anti-CD105/endothelin monoclonal  
14 antibody in astrocytic tumors: Comparison with an anti-CD31 monoclonal  
15 antibody. *Neuropathology*. 25, 201–206.

16 Zhang, Z.G., Zhang, L., Jiang, Q., Zhang, R., Davies, K., Powers, C., Van Bruggen,  
17 N., Chopp, M., 2000. VEGF enhances angiogenesis and promotes blood-  
18 brain barrier leakage in the ischemic brain. *Journal of Clinical Investigation*.  
19 106, 829-838.

20 Zhang, Z.G., Chopp, M., 2015. Promoting brain remodeling to aid in stroke  
21 recovery. In: *Trends in Molecular Medicine*. Vol. 21, ed.^eds., pp. 543-548.

22 Zhu, X., Xie, M., Wang, K., Zhang, K., Gao, Y., Zhu, L., Zhou, F., 2014. The effect  
23 of puerarin against IL-1 $\beta$ -mediated leukostasis and apoptosis in retinal  
24 capillary endothelial cells (TR-iBRB2). *Mol Vis*. 20, 1815-1823.

25

## 26 **Figure legends.**

27 **Figure 1. Effects of B10 transplantation on angiogenesis in a MCAO rat**

28 **model.** B10 cells were transplanted intravenously 24 h after MCAO. Control MCAO

1 rats received PBS instead of cells. Three and 7 days after MCAO, vessels in the  
2 core and IBZ region were visualized by vWF immunofluorescence staining. Vessel  
3 density was evaluated using ImageJ, as described in the Materials and Methods,  
4 and expressed as percent of total area of a microscopic field at x400 magnification.  
5 Representative vWF immunofluorescence staining photomicrographs of ischemic  
6 core and IBZ cortices of rat brains 3 and 7 days after MCAO are shown in (A) and  
7 (C), respectively. Quantified data of the staining of rat brains 3 (B) and 7 (D) days  
8 after MCAO are presented as average of 5 rats  $\pm$  SD. Statistical significance are  
9 denoted as follows; \* $p < 0.05$  vs same area of PBS (control) rats, # $p < 0.001$  vs  
10 same area of PBS (control) rats,  $^{\dagger}p < 0.001$  vs core area of PBS (control) rats.

11 **Figure 2. Detection of transplanted B10 cells and newly formed vessels in the**  
12 **brains of MCAO rats.** B10 cells were transplanted intravenously 24 h after MCAO.  
13 Control MCAO rats received PBS instead of cells. Three and 7 days after MCAO,  
14 B10 in the core and IBZ region were visualized by human nuclear antigen (HuN)  
15 immunofluorescence staining. Similarly, newly formed vessels were identified by  
16 endoglin immunofluorescence staining. Endoglin immunofluorescence was  
17 evaluated using ImageJ, as described in the Materials and Methods.  
18 Representative HuN immunofluorescence photomicrographs of ischemic core and  
19 IBZ cortices of rat brains, 3 days (C) and 7 days (D) after MCAO are shown, and  
20 quantified data of the staining of 3 days (B) and 7 days (D) after MCAO are  
21 presented in (E) and (F), respectively. The quantified data are presented here as

1 average  $\pm$  SD of 5 rats in a group. Statistical significance are denoted as follows;

2 \* $p < 0.05$  vs same area of PBS (control) rats. Bar = 50  $\mu\text{m}$ .

3 **Figure 3. Effects of B10 transplantation on the mRNA level of angiogenesis**

4 **related factors in a MCAO rat model.** B10 cells were transplanted intravenously

5 24 h after MCAO. Control MCAO rats received PBS instead of cells. Sham animals

6 were undergone all surgical procedure except occlusion of the artery and

7 transplantation. Three days after MCAO, cortical brain tissues from ischemic core,

8 IBZ and contralateral areas were dissected out, total RNA was isolated, reverse

9 transcribed, and mRNA levels of angiogenesis related factors were evaluated by

10 real time PCR. The mRNA levels of target genes were normalized with those of

11 corresponding GAPDH, calculated as fold induction relative to mRNA from cortex

12 of a sham animal, and results are presented as mean  $\pm$  SD of 5 rats. Statistical

13 significance is denoted as follows; \* $p < 0.05$ , vs PBS (control) rats.

14 **Figure 4. Effects of B10 transplantation on HIF-1 $\alpha$  protein expression in**

15 **MCAO rat brains.** B10 cells were transplanted intravenously 24 h after MCAO.

16 Control MCAO rats received PBS instead of cells. Three days after MCAO, cortical

17 brain tissues from ischemic core, IBZ and contralateral areas were dissected out,

18 and levels of HIF-1 $\alpha$  and p300 were evaluated by Western blot analysis using

19 specific antibodies (A).  $\beta$ -Actin was used as a loading control (A). To determine the

20 localization of HIF-1 $\alpha$ , double immunofluorescence staining was done where Iba-1

21 (B) and GFAP (C) were used as astrocyte and macrophage/microglia markers,

22 respectively. Co-localization of HIF-1 $\alpha$  and p300 was determined by double

1 immunofluorescence staining using specific antibodies (D). Bar = 50  $\mu$ m in (B and  
2 C) and 20  $\mu$ m in (D)

3 **Figure 5. Effects of B10 transplantation on VEGF protein expression in**  
4 **MCAO rat brains.** B10 cells were transplanted intravenously 24 h after MCAO.  
5 Control MCAO rats received PBS instead of cells. Three days after MCAO, cortical  
6 brain tissues from ischemic core, IBZ and contralateral areas were dissected out,  
7 and level of VEGF was evaluated by Western blot analysis using specific antibody  
8 (A).  $\beta$ -Actin was used as a loading control (A). To determine the distribution and  
9 morphology of VEGF expressing cells in MCAO, immunostaining with specific  
10 antibody was done. Representative photomicrographs of cortical core, IBZ  
11 contralateral areas are shown in (B). Double immunofluorescence staining was  
12 done to identify VEGF-expressing cells where GFAP was used as astrocyte marker  
13 (C), and Iba-1 as macrophage/microglia marker (D). Bar= 50  $\mu$ m.

14 **Figure 6. Effects of B10 transplantation on TGF $\beta$  protein levels in MCAO rat**  
15 **brains.** B10 cells were transplanted intravenously 24 h after MCAO. Control  
16 MCAO rats received PBS instead of cells. Three days after MCAO, cortical brain  
17 tissues from ischemic core, IBZ and contralateral areas were dissected out, and  
18 level of TGF $\beta$  was evaluated by Western blot analysis using specific antibody (A).  
19  $\beta$ -Actin was used as a loading control (A). The protein expression was further  
20 evaluated by immunofluorescence staining using TGF $\beta$ -specific antibody.  
21 Representative photomicrographs of core, IBZ and contralateral cortical areas of

1 PBS-treated and B10-transplanted MCAO rats are shown in (B). Hoechst was used  
2 for nuclear staining.

3 **Figure 7. Influence of B10 and microglia on the regulation of VEGF**

4 **expression.** B10 and HMO6 cells were treated with inflammatory cytokines for  
5 indicated times. After isolation of total RNA, cDNA was prepared, and VEGF  
6 mRNA levels in B10 (A) and HMO6 (B) were evaluated by real time PCR using  
7 gene specific primers. The mRNA was quantified as fold induction relative to  
8 medium-treated condition of same time points after normalization with  
9 corresponding GAPDH mRNA, and the results are presented as mean  $\pm$  SEM of at  
10 least 3 independent experiments. Statistical significance is denoted as follows; \* $p$ <  
11 0.05, vs medium-treated condition of same time point. (C and D) B10 and HMO6  
12 were co-cultured in 0.5% FBS containing DMEM medium for indicated times. After  
13 isolation of total RNA, cDNA was prepared, and the mRNA levels of IL-1 $\beta$  (C) and  
14 VEGF (D) in B10 and HMO6 was determined by real time PCR. After normalization  
15 with corresponding GAPDH mRNA, the mRNA was quantified relatively using 1  
16 sample of HMO6 normal culture as a calibrator. The data presented here as an  
17 average  $\pm$  SD of 3 independent experiments. Statistical significance is denoted as  
18 follows; \* $p$ < 0.05, vs native culture condition of same time point. (E and F) HMO6  
19 was treated 0.5% FBS containing DMEM for 24 h in native culture, or B10 co-  
20 culture (B10 CC) condition. Then the cell lysate was collected before, or 4 h of  
21 OGD. The levels of HIF1 $\alpha$  and VEGF in HMO6 cell lysates were evaluated by  
22 Western blotting using specific antibodies.  $\beta$ -Actin was used as a loading control. A

1 representative Western blotting data is shown in (E), and its  $\beta$ -Actin-normalized  
2 densitometric analysis is shown in (F).

3 **Figure 8. Effects of B10 transplantation on IL-1 $\beta$  expression in MCAO rat**

4 **brains.** B10 cells were transplanted intravenously 24 h after MCAO. Control  
5 MCAO rats received PBS instead of cells. Three days after MCAO, the distribution  
6 of IL-1 $\beta$ -expressing cells are determined by immunostaining. Representative  
7 photomicrographs of PBS and B10 transplanted rats at low magnification are show  
8 in (A), where cortical core region is demarcated by dashed lines. Higher  
9 magnification of photomicrographs of core and IBZ region are shown in (B).

10 Immunoreactive cells are counted in the core region at X400 magnification, and the  
11 data are presented as average  $\pm$  SD of 5 rats in (C). To determine IL-1 $\beta$   
12 expressing cells in the core, double immunofluorescence staining was done where  
13 ED-1 and human nuclear antigen were used as macrophage/microglia and  
14 transplanted B10 cell marker, respectively. Representative photomicrographs of  
15 double immunofluorescence staining of IL-1 $\beta$  and ED-1 in PBS and B10  
16 transplanted rats are shown in (D). Double immunofluorescence staining of IL-1 $\beta$   
17 and human nuclear antigen in PBS and B10 transplanted rats are shown in (E).  
18 Statistical significance is denoted as follows; \* $p$  < 0.05, vs PBS (control) rats. Bar=  
19 100  $\mu$ m in (A), and 50  $\mu$ m in (B).

20 **Figure 9. Effects of IL-1 $\beta$ -silenced B10 transplantation on the newly formed**

21 **vessels in MCAO rat brains.** IL-1 $\beta$ -silenced B10, or non-silenced B10 cells were  
22 transplanted intravenously 24 h after MCAO. Three and 7 days after MCAO, newly

1 formed vessels in the core and IBZ region were visualized by endoglin  
2 immunofluorescence staining. The density of newly formed vessels was evaluated  
3 using ImageJ, as described in the Materials and Methods, and expressed as  
4 percent of total area of a microscopic field at x400 magnification. Representative  
5 endoglin immunofluorescence staining photomicrographs of ischemic core and IBZ  
6 cortices of rat brains 3 and 7 days after MCAO are shown in (A) and (C),  
7 respectively. Boxed area of the merged photomicrographs are enlarged and shown  
8 in 4<sup>th</sup> column. Vessel-like structures are indicated by arrows. Quantified data of the  
9 staining of rat brains 3 (B) and 7 (D) days after MCAO are presented as average of  
10 5 rats  $\pm$  SD. Statistical significance are denoted as follows; \* $p$ < 0.01 vs same area  
11 of non-silenced B10 transplanted MCAO rats.

12

### 13 **Supplemental Methods:**

14 A human neuronal cell line (A1) was generated by somatic fusion of a human  
15 primary fetal cerebral neuron and a human neuroblastoma cell (Nagai et al., 2002).  
16 A1 cells showed similar morphological, electrophysiological and expressional  
17 features like primary neurons in culture. A1 cells were cultured in 5% FBS  
18 containing DMEM medium. For differentiation, 10  $\mu$ M of retinoic acid was used for  
19 48 h.

### 20 **Supplemental reference:**

21 Nagai, A., Suzuki, Y., Baek, S.Y., Lee, K.S., Lee, M.C., McLarnon, J.G., Kim, S.U.,  
22 2002. Generation and characterization of human hybrid neurons produced  
23 between embryonic CNS neurons and neuroblastoma cells. *Neurobiol Dis.* 11,184-  
24 198

25

1 **Supplemental figure legends**

2 **Supplemental figure 1. Effects of B10 transplantation on vessel density in a**  
3 **14-day old MCAO rat model.** B10 cells were transplanted intravenously 24 h after  
4 MCAO. Control MCAO rats received PBS instead of cells. Fourteen days after  
5 MCAO, vessels in the core and IBZ region were visualized by vWF  
6 immunofluorescence staining. Representative vWF immunofluorescence staining  
7 photomicrographs of ischemic core and IBZ cortices of rat brains are shown here.  
8 Bar = 50  $\mu$ m.

9 **Supplemental figure 2. Effects of B10 transplantation on Endoglin<sup>+</sup> newly**  
10 **formed vessel density in a 14-day old MCAO rat model.** B10 cells were  
11 transplanted intravenously 24 h after MCAO. Control MCAO rats received PBS  
12 instead of cells. Fourteen days after MCAO, newly formed vessels in the core and  
13 IBZ region were visualized by Endoglin immunofluorescence staining.  
14 Representative Endoglin immunofluorescence staining photomicrographs of  
15 ischemic core and IBZ cortices of rat brains are shown here. Bar = 50  $\mu$ m.

16 **Supplemental figure 3. Expression of IL-1 $\beta$  protein B10 (mesenchymal stem**  
17 **cell line), HMO6 (microglia cell line) and A1 (neuronal hybridoma cell line)**  
18 **cells.** B10, HMO6 and A1 cells were cultured until about 80% confluency. Then the  
19 cells were cultured in 0.5% FBS containing DMEM for 24 h. The culture  
20 supernatant was collected and subjected to ELISA to quantify IL-1 $\beta$ . The data  
21 presented here as average  $\pm$  SD of 3 independent experiments.

1 **Supplemental figure 4: Silencing IL-1 $\beta$  expression in B10 cells by**  
2 **transfecting IL-1 $\beta$ -specific siRNA.** To silence IL-1 $\beta$  expression, mRNA-specific  
3 silencing siRNA was transfected in B10 cells, as described in the Materials and  
4 Methods. Forty eight hours after transfection, the cells were cultured in 0.5% FBS  
5 containing DMEM for 24 h. Total RNA was isolated, reverse transcribed and IL-1 $\beta$   
6 mRNA levels were evaluated by real time PCR. The level of IL-1 $\beta$  mRNA was  
7 calculated by relative quantification method using a RNA sample of non-  
8 transfected B10 culture as calibrator, and shown in (A). (B) In another experiment,  
9 48 h after transfection, B10 cells were cultured in 0.5% FBS containing DMEM for  
10 24 h. Then the culture supernatant and the cell lysate were collected. The level of  
11 secreted IL-1 $\beta$  in culture supernatant was evaluated by ELISA, and the data was  
12 normalized with total protein content of the lysate. ELISA data are shown here  
13 as % control, where a non-transfected B10 culture was served as control. The data  
14 presented here as average  $\pm$  SD of 3 independent experiments. Statistical  
15 significance is denoted as \* $p$  <0.05 vs mock transfected samples.